

SATAKE-FURSTENBERG COMPACTIFICATIONS, THE MOMENT MAP AND λ_1

LEONARDO BILIOTTI AND ALESSANDRO GHIGI

ABSTRACT. Let G be a complex semisimple Lie group, K a maximal compact subgroup and τ an irreducible representation of K on V . Denote by M the unique closed orbit of G in $\mathbb{P}(V)$ and by $\mathcal{O}$ its image via the moment map. For any measure γ on M we construct a map Ψ_γ from the Satake compactification of G/K (associated to V) to the Lie algebra of K . If γ is the K -invariant measure, then Ψ_γ is a homeomorphism of the Satake compactification onto the convex envelope of $\mathcal{O}$. For a large class of measures the image of Ψ_γ is the convex envelope. As an application we get sharp upper bounds for the first eigenvalue of the Laplacian on functions for an arbitrary Kähler metric on a Hermitian symmetric space.

CONTENTS

1. Introduction	1
2. Flag manifolds and compactifications	4
2.1. Flag manifolds	4
2.2. Satake compactifications	6
2.3. τ -connected subspaces	8
2.4. Projections and rational maps	11
2.5. τ -connected subspaces and the moment map	14
3. The Bourguignon-Li-Yau map	19
3.1. Definition and boundary behaviour	19
3.2. The case of the K -invariant measure	23
3.3. Furstenberg compactifications	25
4. Application to eigenvalue estimates	26
References	30

1. INTRODUCTION

Let M be a closed manifold of dimension n and let $\mathcal{M}(M)$ be the set of Riemannian metrics on M . For $g \in \mathcal{M}(M)$ denote by $\lambda_1(M, g)$ the first eigenvalue of $-\Delta_g$, where Δ_g is seminegative the Laplace-Beltrami operator on functions. The

2000 *Mathematics Subject Classification.* 32M10; 53C35; 58J50.

The first author was supported by the Project MIUR “Geometric Properties of Real and Complex Manifolds” and by GNSAGA of INdAM. The second author was supported by the PRIN 2007 MIUR “Moduli, strutture geometriche e loro applicazioni” and by GNSAGA of INdAM. .

quantity $\lambda_1(M, g) \operatorname{vol}(M, g)^{2/n}$ is scale invariant. In 1970 Hersch [20] proved that

$$\sup_{g \in \mathcal{M}(S^2)} \lambda_1(S^2, g) \operatorname{vol}(S^2, g) = 8\pi.$$

A similar inequality was proved by Yang and Yau [37] on higher genus surfaces. On the other hand Colbois and Dodziuk [11] proved that if $n \geq 3$, the quantity $\lambda_1(M, g) \operatorname{vol}(M, g)^{2/n}$ is not bounded above (see also [6], [8] and [12]). If M is a Kähler manifold and $a \in H^2(M)$, denote by $\mathcal{K}(a)$ the set of Kähler metrics with Kähler form in a . It makes sense to study

$$(1) \quad I(a) := \sup_{g \in \mathcal{K}(a)} \lambda_1(M, g).$$

In 1994 Bourguignon, Li and Yau proved that $I(a) < +\infty$ for a large class of pairs (M, a) (including all projective ones) and gave an estimate of $I(a)$ in terms of holomorphic maps $M \rightarrow \mathbb{P}^n$. The main tool in their proof is an interesting geometric construction relating the symmetric space $\operatorname{SL}(n+1, \mathbb{C})/\operatorname{SU}(n+1)$ to the set of positive definite Hermitian matrices with trace 1. By the spectral theorem the latter set is the convex hull of the image of the “Veronese embedding”, that is the map which sends a line $\ell \in \mathbb{P}^n$ to the orthogonal projection on ℓ . If one uses the obvious isomorphism to substitute Hermitian matrices with trace 1 with tracefree skew-Hermitian matrices, the “Veronese embedding” becomes just the moment map. This interpretation was already used in a different direction in [2]. Hence the construction of Bourguignon, Li and Yau relates $\operatorname{SL}(n+1, \mathbb{C})/\operatorname{SU}(n+1)$ to the convex hull of the coadjoint orbit corresponding to $\mathbb{P}^n$.

The present paper has two purposes: 1) generalize this construction by replacing $\mathbb{P}^n$ with an arbitrary flag manifold (i.e. a homogeneous space G/P with G a complex semisimple Lie group and P a parabolic subgroup); 2) get sharp upper bounds for λ_1 on any Hermitian symmetric space.

The generalization (1) is as follows. Let $M = G/P$ be a flag manifold. The space $\operatorname{SL}(n+1, \mathbb{C})/\operatorname{SU}(n+1)$ is replaced by the symmetric space $X = G/K$, where K is a maximal compact subgroup. Let ω be a K -invariant integral Kähler form, $\Phi : M \rightarrow \mathfrak{k}^*$ the moment map, $\mathcal{O} = \Phi(M)$ the corresponding coadjoint orbit and denote by $\widehat{\mathcal{O}}$ the convex hull of $\mathcal{O}$. (M, ω) is the unique closed orbit of G in $\mathbb{P}(V)$ for some irreducible representation $\tau : G \rightarrow \operatorname{GL}(V)$. Using τ one can construct a Satake compactification $\overline{X}_\tau^S$ of X . Given a probability measure γ on M we consider the *Bourguignon-Li-Yau map*

$$\Psi_\gamma : X \rightarrow \mathfrak{k}^* \quad \Psi_\gamma(gK) = \int_M \Phi \left(\sqrt{g\theta(g^{-1})} \cdot x \right) d\gamma(x)$$

(θ denotes the Cartan involution; see p. 13 for the square root). We say that γ is τ -admissible if it does not charge the hyperplane sections of $M \subset \mathbb{P}(V)$. Our results are the following.

Theorem 2. *The map Ψ_γ extends to $\overline{X}_\tau^S$. If μ is the K -invariant probability measure on M , then Ψ_μ is a homeomorphism of $\overline{X}_\tau^S$ onto $\widehat{\mathcal{O}}$.*

Both $\overline{X}_\tau^S$ and $\widehat{\mathcal{O}}$ have a rich boundary structure, given respectively by boundary components in the sense of Satake and by open faces in the sense of convex geometry. The homeomorphism Ψ_μ preserves these structures, sending each boundary component diffeomorphically onto an open face.

Theorem 3. *If γ is a τ -admissible measure, then $\Psi_\gamma(\overline{X}_\tau^S) = \widehat{\mathcal{O}}$ and $\Psi_\gamma(\partial\overline{X}_\tau^S) \subset \partial\widehat{\mathcal{O}}$.*

In particular a Satake compactification of a symmetric space G/K of Cartan type IV (i.e. $G = K^\mathbb{C}$) is homeomorphic to the convex body $\Omega := \text{int } \widehat{\mathcal{O}}$ in $\mathfrak{k}$. This fact is not new: Korányi [26] recently showed that for any symmetric space not necessarily of type IV a Satake compactification $\overline{X}_\tau^S$ is homeomorphic to a convex body in $\mathfrak{p}$ where $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. Already for $\mathbb{P}^1$ the map used by Korányi is different from Ψ_μ and the proofs are completely different as well. Our main tool is the moment map and the main point is the link between $\overline{X}_\tau^S$ and the flag manifold M , as described below.

It would be interesting to generalize this construction to Satake compactifications of symmetric spaces not necessarily of type IV, by replacing flag manifolds by real flag manifolds. We leave this to further inquiry.

Following the strategy of Bourguignon, Li and Yau we apply Theorem 3 to get the following result.

Theorem 4. *If M is a Hermitian symmetric space of the compact type, then $I(2\pi c_1(M)) = 2$. The bound is attained by the symmetric metric.*

This bound was previously known in the following special cases: $M = S^2$, proven by Hersch [20], $M = \mathbb{P}^n$ proven by Bourguignon, Li and Yau [10], M the complex Grassmannian, proven by Arezzo, Loi and second author [2], M an irreducible symmetric space whose automorphism group is a classical group, proven in [7]. We remark that in the statement above M can be reducible. It would be interesting to know if this bound holds more generally for any flag manifold. Our proof breaks down since in such generality Lemma 93 is false and the form α is not even closed.

We now describe the contents of the paper. Consider a set of data $G, K, \tau, \langle \cdot, \cdot \rangle$, where G is a connected semisimple complex Lie group, K is a maximal compact subgroup, $\tau : G \rightarrow \text{GL}(V)$ is an irreducible representation with finite kernel and $\langle \cdot, \cdot \rangle$ is a K -invariant Hermitian product on V . Out of these data one can construct a flag manifold on one side and a Satake compactification $\overline{X}_\tau^S$ of $X = G/K$ on the other. This is recalled with some detail in §§2.1-2.2. Next (§2.3) we study the relation between the compactification $\overline{X}_\tau^S$ and the flag manifold M . The boundary components of $\overline{X}_\tau^S$ have been described in terms of root data in the pioneering work of Satake [31]. We recast Satake's analysis in more geometric terms. In particular μ_τ -connected subsets of the simple root are replaced by τ -connected subspaces $W \subset V$, a class of subspaces that are particularly well adapted to M , e.g. the intersection $M_W := M \cap \mathbb{P}(W)$ is a smaller flag manifold. By means of the moment map we show that the submanifolds M_W capture much of the information contained in W . Eventually $\overline{X}_\tau^S$ can be embedded in the set of rational self-maps of M and the boundary component corresponding to M_W corresponds to the rational maps $M \dashrightarrow M_W$ (§2.4). §2.5 is probably the most technical part. Using the Atiyah-Guillemin-Sternberg convexity theorem and a computation in terms of root data, we show that each τ -connected subspace W determines a subset $C_W \subset \mathfrak{k}^*$ defined by linear inequalities, see (54), such that $\mathcal{O} \subset C_W$. If E_W denotes the set where these inequalities become equalities, then $\mathcal{O} \cap E_W = \Phi(M_W)$ (Theorem 57). In words, each τ -connected subspace W gives rise to some inequalities satisfied by $\widehat{\mathcal{O}}$ and is responsible for a part of its boundary. The Bourguignon-Li-Yau map is

defined in §3.1. The interpretation of $\overline{X}_\tau^S$ in terms of rational maps immediately yields the extension of Ψ_γ to $\overline{X}_\tau^S$, while the results of §2.5 allow to prove that if γ is τ -admissible, then Ψ_γ maps the boundary of $\overline{X}_\tau^S$ to the boundary of $\widehat{\mathcal{O}}$. In 3.2 we show that for $\gamma = \mu$ (i.e. the K -invariant measure) the restriction of Ψ_μ to X is a local diffeomorphism. This finally allows to complete the proofs of Theorems 2 and 3. As a byproduct of Theorem 2 we also get a short proof (only for type IV) of a theorem by Moore [28], stating that Satake and Furstenberg compactifications coincide (§3.3). In §4 we give the application to λ_1 and prove Theorem 4.

After completing this work we became aware of preprint [30] by Sanyal, Sottile and Sturmfels, which is devoted to the study of *orbitopes*, which are by definition the convex hulls of orbits of a compact group acting linearly on a real vector space. Our $\widehat{\mathcal{O}}$ is an example of orbitope in the adjoint representation. Although there is no real overlap between [30] and the present paper and although the points of view are rather different, we believe that their more general approach should in the future shed some light on our constructions and conversely the approach in this paper should be of interest also for more general orbitopes.

Acknowledgements. We wish to thank Laura Geatti, Peter Heinzner, Lizhen Ji, Karl-Hermann Neeb, Giorgio Ottaviani and Luiz San Martin for interesting emails/discussions.

2. FLAG MANIFOLDS AND COMPACTIFICATIONS

2.1. Flag manifolds. Let G be a connected complex semisimple Lie group and $K \subset G$ a maximal compact subgroup. Denote by $\theta : G \rightarrow G$ the Cartan involution, such that $K = \text{Fix}(\theta)$. Let $\mathfrak{g}$ and $\mathfrak{k}$ be the Lie algebras and let θ denote also the involution on $\mathfrak{g}$. Consider an irreducible complex representation $\tau : G \rightarrow \text{SL}(V)$ and a Hermitian product $\langle \cdot, \cdot \rangle$ on V , which is invariant by K . This means that for any $g \in G$

$$(5) \quad \tau(\theta(g)) = (\tau(g^{-1}))^*.$$

We will always assume that $\ker \tau$ is finite, i.e. that τ is nontrivial on any simple factor of G .

G, K, τ and $\langle \cdot, \cdot \rangle$ are the basic data for the construction of two different objects: a G -homogeneous complex manifold with a K -invariant Hodge metric on the one side and a compactification of the space $X = G/K$ on the other side. The first one is a flag manifold, the second is a Satake compactification. In this § we describe the flag manifold. The definition of the Satake compactification will be recalled in §2.2.

Remark 6. *If we compose τ with the natural map $\text{SL}(V) \rightarrow \mathbb{P}\text{GL}(V)$ and factor by the (finite) kernel of the composition, we get a faithful projective representation of some finite quotient of G . Conversely, any faithful representation of G can be lifted to a linear representation of some finite covering of G . Strictly speaking the basic object in the construction of flag manifolds and Satake compactifications is the projective representation, see [31, p. 85] and [9, p. 63]. Nevertheless both constructions are not affected by passing to finite quotients/coverings (compare [18, §4.1]). For simplicity we will deal with the representation $\tau : G \rightarrow \text{SL}(V)$.*

The K -invariant Hermitian product on V gives rise to a K -invariant Kähler metric on $\mathbb{P}(V)$, namely the Fubini-Study metric associated to $\langle \cdot, \cdot \rangle$. Since τ is irreducible, in $\mathbb{P}(V)$ there is exactly one closed orbit of G , which coincides with the unique complex orbit of K (see [21, p. 124]). We denote this orbit by M and endow it with the restriction ω of the Fubini-Study metric. Thus (M, ω) is a Kähler manifold and K acts transitively, symplectically and almost effectively on M . Let $\xi_v \in \mathfrak{X}(M)$ denote the fundamental vector field corresponding to $v \in \mathfrak{k}$. Since M is simply connected and K is connected and semisimple the action of K is Hamiltonian with a unique moment map $\Phi : M \rightarrow \mathfrak{k}^*$ (see e.g. [17, Thm. 26.1 pp. 185-187]), that for any $v \in \mathfrak{k}$ and any $a \in K$ satisfies

$$(7) \quad d\langle \Phi, v \rangle = -i_{\xi_v} \omega \quad \Phi(ax) = \text{Ad}^*(a)(\Phi(x)) = \Phi(x) \circ \text{Ad}(a^{-1}).$$

Φ is a diffeomorphism of M onto a coadjoint orbit of K that we denote by $\mathcal{O} := \Phi(M)$ (see e.g. [17, Thm. 32.17 p. 260]). We call M , endowed with all these structures, the *flag manifold* associated to G, K, τ and $\langle \cdot, \cdot \rangle$.

Denote by B the Killing form of $\mathfrak{g}$ and by

$$\langle \cdot, \cdot \rangle : \mathfrak{k}^* \times \mathfrak{k} \rightarrow \mathbb{R}$$

the duality pairing. Fix a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ such that $\theta(\mathfrak{h}) = \mathfrak{h}$. Set

$$(8) \quad \mathfrak{t} := \mathfrak{h} \cap \mathfrak{k} \quad \mathfrak{a} := i\mathfrak{t}.$$

$\mathfrak{t}$ is the Lie algebra of a maximal torus $T \subset K$ and $\mathfrak{h} = \mathfrak{t} \oplus \mathfrak{a}$. Denote by $\Delta(\mathfrak{g}, \mathfrak{h})$ the root system of $(\mathfrak{g}, \mathfrak{h})$. Let Π be a system of simple roots for $(\mathfrak{g}, \mathfrak{h})$. Then we say that $(\mathfrak{h}, \Pi)$ is a *root datum*. (By this we also understand that $\mathfrak{h}$ is θ -stable.) We denote by $\Delta_+ = \Delta_+(\mathfrak{h}, \Pi)$ the corresponding set of positive roots and put $\Delta_- = -\Delta_+$. For $\alpha \in \Delta$ let $H_\alpha \in \mathfrak{h}$ be such that $\alpha(X) = B(X, H_\alpha)$ for any $X \in \mathfrak{h}$. Also we transfer $B|_{\mathfrak{h} \times \mathfrak{h}}$ to a bilinear form on $\mathfrak{h}^*$ by the definition

$$(9) \quad B(\alpha, \beta) = B(H_\alpha, H_\beta) = \alpha(H_\beta) = \beta(H_\alpha)$$

for $\alpha, \beta \in \mathfrak{h}^*$. Denote by $\mathfrak{b}_+(\mathfrak{h}, \Pi)$ the standard *positive* Borel subalgebra:

$$(10) \quad \mathfrak{b}_+(\mathfrak{h}, \Pi) = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta_+} \mathfrak{g}_\alpha$$

Given a set $I \subset \Pi$ of simple roots put

$$(11) \quad \begin{aligned} \Delta_I &= \Delta \cap \text{span}(I) & \mathfrak{p}_I &= \mathfrak{b}_+(\mathfrak{h}, \Pi) \oplus \bigoplus_{\alpha \in \Delta_I \cap \Delta_-} \mathfrak{g}_\alpha \\ \mathfrak{h}_I &= \bigcap_{\alpha \in I} \ker \alpha & \mathfrak{h}^I &= \bigoplus_{\alpha \in I} \mathbb{C}H_\alpha \\ \mathfrak{s}_I &= \mathfrak{h}^I \oplus \bigoplus_{\alpha \in \Delta_I} \mathfrak{g}_\alpha & \mathfrak{u}_I &= \bigoplus_{\alpha \in \Delta_+ - \Delta_I} \mathfrak{g}_\alpha. \end{aligned}$$

(E.g. $\mathfrak{p}_\emptyset = \mathfrak{b}_+(\mathfrak{h}, \Pi)$ and $\mathfrak{p}_\Pi = \mathfrak{g}$.) Then $\mathfrak{p}_I$ is a parabolic subalgebra of $\mathfrak{g}$ and any parabolic subalgebra containing $\mathfrak{b}_+(\mathfrak{h}, \Pi)$ is of this form. There is a decomposition $\mathfrak{h} = \mathfrak{h}_I \oplus \mathfrak{h}^I$ which is B -orthogonal. $\mathfrak{u}_I$ is a nilpotent ideal of $\mathfrak{p}_I$, while $\mathfrak{s}_I$ is a semisimple subalgebra of $\mathfrak{g}$ which commutes with $\mathfrak{h}_I$. Denote by P_I, U_I and S_I the connected subgroups of G with Lie algebras $\mathfrak{p}_I, \mathfrak{u}_I$ and $\mathfrak{s}_I$ respectively. They are closed subgroups, P_I is parabolic, U_I is the unipotent radical of P_I , while S_I is semisimple. Moreover $\mathfrak{h}_I \oplus \mathfrak{s}_I$ is a reductive subalgebra and one has the *Chevalley* (or *algebraic Levi*) *decomposition* $\mathfrak{p}_I = \mathfrak{u}_I \oplus \mathfrak{h}_I \oplus \mathfrak{s}_I$ (see e.g. [34, p. 32]).

Once a root datum $(\mathfrak{h}, \Pi)$ has been fixed, the representation τ determines the line $x_0 = \mathbb{C}v_\tau$ spanned by any highest weight vector v_τ and $M = G \cdot x_0 = K \cdot x_0$. The stabilizer of x_0 in G is a parabolic subgroup P which does not contain any simple factor of G , and $K_0 = K \cap P$ is the centralizer of a subtorus of T [33, Thm. 1]. The following computation is well-known (see e.g. [38] or [5, p.63], that has the opposite sign convention for Φ). We recall the proof for the reader's convenience.

Proposition 12. *Let μ_τ be the highest weight of τ and let v_τ be a highest weight vector. Set $x_0 = [v_\tau]$ and for any $X \in \mathfrak{g}$ write $X = X^\mathfrak{h} + \sum_{\alpha \in \Delta} X_\alpha$ where $X^\mathfrak{h} \in \mathfrak{h}$ and $X_\alpha \in \mathfrak{g}_\alpha$. Then*

$$(13) \quad \langle \Phi(x_0), X \rangle = -i\mu_\tau(X^\mathfrak{h})$$

Proof. Fix on $\mathbb{P}(V)$ the Fubini-Study metric induced by $\langle \cdot, \cdot \rangle$. The moment map $\Phi^{\mathbb{P}(V)}$ of $(\mathbb{P}(V), \omega_{FS})$ with respect to the $SU(V)$ -action is given by the formula

$$\langle \Phi^{\mathbb{P}(V)}([v]), A \rangle = -i \frac{\langle Av, v \rangle}{|v|^2} \quad A \in \mathfrak{su}(V), \quad v \in V.$$

Since the inclusion $M \hookrightarrow \mathbb{P}(V)$ is K -equivariant and symplectic $\Phi^M(x) = \Phi^{\mathbb{P}(V)}(x) \circ \tau$ where $\tau : \mathfrak{k} \rightarrow \mathfrak{su}(V)$ denotes the infinitesimal representation. So for any $X \in \mathfrak{k}$

$$\langle \Phi^M(x_0), X \rangle = -i \frac{\langle \tau(X)v_\tau, v_\tau \rangle}{|v_\tau|^2}.$$

Write $X = X^\mathfrak{h} + \sum_\alpha X_\alpha$ with $X_\alpha \in \mathfrak{g}_\alpha$. Since $\tau(X_\alpha)v_\tau \in V_{\mu_\tau + \alpha} \perp V_{\mu_\tau}$

$$\langle \tau(X)v_\tau, v_\tau \rangle = \langle \tau(X^\mathfrak{h})v_\tau, v_\tau \rangle = \mu_\tau(X^\mathfrak{h})|v_\tau|^2.$$

This yields the result. $\square$

2.2. Satake compactifications. Assume given a *real* semisimple Lie group G , a maximal compact subgroup $K \subset G$ and an infinitesimally faithful irreducible representation τ of G on a complex vector space V , which is endowed with a K -invariant Hermitian product. With these data Satake [31] constructed a compactification $\overline{X}_\tau^S$ of the symmetric space $X = G/K$. In this paper we are concerned only with the case in which $G = K^\mathbb{C}$ (i.e. X is of Cartan type IV). In this case the Lie theoretic data simplify considerably since the restricted roots of the real Lie algebra $\mathfrak{g}_\mathbb{R}$ underlying $\mathfrak{g}$ coincide with the roots of the complex Lie algebra $\mathfrak{g}$. We wish to recall the construction of the Satake compactifications and some of their relevant properties restricting to this particular class of symmetric spaces and taking advantage of this simplification. Proofs can be found in the general case in the book [9, §I.1] which we follow for most of the notation.

Put

$$\mathcal{H}(V) = \{A \in \text{End}(V) : A = A^*\}.$$

Denote by $\pi : \mathcal{H}(V) \setminus \{0\} \rightarrow \mathbb{P}(\mathcal{H}(V))$ the canonical projection and set

$$\mathcal{P}(V) = \pi(\{A \in \mathcal{H}(V) : A > 0\}) \subset \mathbb{P}(\mathcal{H}(V)).$$

$\mathcal{P}(V)$ consists of points $[A]$ such that A is invertible and all its eigenvalues have the same sign.

Lemma 14. (a) $\overline{\mathcal{P}(V)} = \pi(\{A \in \mathcal{H}(V) : A \neq 0, A \geq 0\})$. (b) The restriction of π to $\{A \in \mathcal{H}(V) : A > 0, \det A = 1\}$ is a homeomorphism onto $\mathcal{P}(V)$. (c) The restriction of π to $\{A \in \mathcal{H}(V) : A \geq 0, \text{tr } A = 1\}$ is a homeomorphism onto $\overline{\mathcal{P}(V)}$.

Definition 15. For $G, K, \tau, \langle \cdot, \cdot \rangle$ as at p. 4 set

$$(16) \quad i_\tau : X := G/K \rightarrow \mathcal{P}(V) \quad i_\tau(gK) = [\tau(g)\tau(g)^*].$$

The Satake compactification of X associated to τ and $\langle \cdot, \cdot \rangle$ is the space $\overline{X}_\tau^S := \overline{i_\tau(X)}$. The closure is taken in $\mathbb{P}(\mathcal{H}(V))$.

Since $\mathrm{SL}(V)$ and hence G act on $\mathbb{P}(\mathcal{H}(V))$ by conjugation, $\overline{X}_\tau^S$ is a G -compactification. We stress that $\overline{X}_\tau^S$ depends only on G, K, τ and $\langle \cdot, \cdot \rangle$.

Satake gave a thorough description of the boundary $\partial \overline{X}_\tau^S := \overline{X}_\tau^S - i_\tau(X)$ in terms of root data. We will now recall this description in our simplified setting. In §2.3 we will reinterpret this description in a way that does not depend on the root data.

Definition 17. Fix a root datum $(\mathfrak{h}, \Pi)$. A subset $E \subset \mathfrak{a}^*$ is connected if there is no pair of disjoint subsets $D, C \subset E$ such that $D \cup C = E$, and $B(\lambda, \mu) = 0$ for any $\lambda \in D$ and $\mu \in C$.

(A thorough discussion of connected subsets can be found in [28, §5].) Connected components are defined as usual. For example the connected components of Π are the subsets corresponding to the simple roots of the simple ideals in $\mathfrak{g}$. Denote by μ_τ the highest weight of τ with respect to $(\mathfrak{h}, \Pi)$ and let v_τ be a highest weight vector.

Definition 18. A subset $I \subset \Pi$ is μ_τ -connected if $I \cup \{\mu_\tau\}$ is connected.

Equivalently, I is μ_τ -connected if and only if every connected component of I contains at least one element α with $B(\mu_\tau, \alpha) = \mu_\tau(H_\alpha) \neq 0$. By the highest weight theorem, if $\lambda \in \mathfrak{a}^*$ is a weight of τ , then

$$\lambda = \mu_\tau - \sum_{\alpha \in \Pi} c_\alpha \alpha$$

for some nonnegative integers c_α . The support of λ is the set $\mathrm{supp}(\lambda) := \{\alpha \in \Pi : c_\alpha > 0\}$.

Lemma 19 ([31, Lemma 5 p. 87]). $I \subset \Pi$ is μ_τ -connected if and only if $I = \mathrm{supp}(\lambda)$ for some weight λ of τ .

For example $\emptyset = \mathrm{supp}(\mu_\tau)$ and Π is μ_τ -connected since τ is nontrivial on any simple factor of G . Given $\lambda \in \mathfrak{h}^*$ denote by V_λ the corresponding eigenspace and set

$$V_I = \bigoplus_{\mathrm{supp}(\lambda) \subset I} V_\lambda.$$

Lemma 20 ([31, Lemma 8 p. 89]). Let S_I be the subgroup of G defined in (11). Then $\tau(g)(V_I) \subset V_I$ for any $g \in S_I$ and the representation $\tau_I : S_I \rightarrow \mathrm{GL}(V_I)$ gotten in this way is irreducible. The highest weight of τ_I is $\mu_\tau|_{\mathfrak{h}_I}$ and $v_\tau \in V_I$ is a highest weight vector.

Definition 21. If $I \subset \Pi$ is μ_τ -connected, denote by I' the collection of all simple roots orthogonal to $\{\mu_\tau\} \cup I$. The set $J := I \cup I'$ is called the μ_τ -saturation of I .

The largest μ_τ -connected subset contained in J is I .

Lemma 22 ([9, Prop. I.4.29 p. 70]). *If I is μ_τ -connected, then $P_J = \{g \in G : \tau(g)V_I \subset V_I\}$.*

Fix a μ_τ -connected subset I . If $A \in \text{End}(V_I)$, let $A \oplus 0$ denote the extension of A that is trivial on $V_I^\perp$. If $\pi_I : V \rightarrow V_I$ denotes orthogonal projection and $j_I : V_I \hookrightarrow V$ denotes the inclusion, then $A \oplus 0 = j_I \circ A \circ \pi_I$ and the map

$$(23) \quad \psi : \mathbb{P}(\mathcal{H}(V_I)) \rightarrow \mathbb{P}(\mathcal{H}(V)) \quad \psi([A]) = ([A \oplus 0])$$

embeds $\mathcal{P}(V_I)$ in $\overline{\mathcal{P}(V)}$. Note that $K_I = S_I \cap K$ is a maximal compact subgroup of S_I and S_I is a semisimple complex Lie group. Therefore $X_I = S_I/K_I$ is again a symmetric space of type IV. Hence we have a map $i_{\tau_I} : X_I \rightarrow \mathcal{P}(V_I)$ defined as in (16). Finally define

$$i_I = \psi \circ i_{\tau_I} : X_I \rightarrow \overline{\mathcal{P}(V)}.$$

Theorem 24 ([9, Cor. I.4.32]).

$$\overline{X}_\tau^S = \bigsqcup_{\mu_\tau\text{-connected } I} G \cdot i_I(X_I).$$

If $I = \Pi$ then $i_I(X_I) = X$. The sets $g \cdot i_I(X_I)$ with $g \in G$ and $I \subsetneq \Pi$ are called *boundary components*.

Lemma 25 ([9, Prop. I.4.29]). *The boundary components are disjoint: if I is μ_τ -connected and $g \in G$ then $g \cdot i_I(X_I) \cap i_I(X_I) \neq \emptyset$ if and only if $g \cdot i_I(X_I) = i_I(X_I)$ if and only if $g \in P_J$.*

2.3. τ -connected subspaces. We wish to interpret the construction of Satake compactifications more intrinsically, i.e. independently of the root data. Given a subspace $W \subset V$, $W \neq \{0\}$, set

$$P_W = \{g \in G : \tau(g)W \subset W\} \quad \tilde{K}_W = K \cap P_W \quad M_W = M \cap \mathbb{P}(W).$$

The subgroup P_W is closed.

Definition 26. *W is a τ -connected subspace if P_W is parabolic and acts irreducibly on W .*

Lemma 27. *If W is τ -connected, then $\tilde{K}_W$ is connected and contains a maximal torus of K . There is a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ such that $\theta(\mathfrak{h}) = \mathfrak{h}$ and $\mathfrak{h} \subset \mathfrak{p}_W$.*

Proof. Since $G/P_W = K/\tilde{K}_W$ by [33, Thm. 1] $\tilde{K}_W$ is the centralizer of a torus contained in K . Therefore it is connected and contains a maximal torus T . If $\mathfrak{t} = \text{Lie } T$, then $\mathfrak{h} := \mathfrak{t} \oplus i\mathfrak{t}$ is a Cartan subalgebra with the required properties. $\square$

Corollary 28. *For any τ -connected subspace W there are a root datum $(\mathfrak{h}, \Pi)$ and a subset $J \subset \Pi$ such that*

$$(29) \quad \theta(\mathfrak{h}) = \mathfrak{h} \quad \mathfrak{b}_+(\mathfrak{h}, \Pi) \subset \mathfrak{p}_W \quad \mathfrak{p}_W = \mathfrak{p}_J.$$

($\mathfrak{p}_J$ is as in (11).) *There are many choices for $(\mathfrak{h}, \Pi)$. If one choice is fixed, then J is unique.*

Denote by R_W the Zariski closure of $\tilde{K}_W$. It is a complex connected subgroup of G with Lie algebra $\mathfrak{r}_W = \tilde{\mathfrak{k}}_W \otimes \mathbb{C}$. Let $\tilde{S}_W = (R_W, R_W)$ be the commutator subgroup of R_W . It is a connected complex semisimple Lie group. Therefore it splits as a product of simple factors $\tilde{S}_W = S_1 \times \cdots \times S_r$. We can reorder the factors

in such a way that the action of $S_1, \dots, S_q$ on W be nontrivial, while the remaining factors $S_{q+1}, \dots, S_r$ act trivially on W . Set

$$(30) \quad \begin{aligned} S_W &:= S_1 \times \cdots \times S_q & S'_W &:= S_{q+1} \times \cdots \times S_r \\ K_W &= S_W \cap K & K'_W &= S'_W \cap K. \end{aligned}$$

S_W and S'_W are closed complex connected semisimple subgroups of G , while K_W and K'_W are maximal compact subgroups of S_W and S'_W respectively. Finally denote by $\tilde{\tau}_W$ and τ_W the representations of $\tilde{S}_W$ and S_W on W induced by τ . We stress that all these definitions depend only on W .

Proposition 31. *Let $W \subset V$ be a τ -connected subspace and let $(\mathfrak{h}, \Pi)$ and J be as in (29). Then*

- a) $\mathfrak{r}_W = \mathfrak{h}_J \oplus \mathfrak{s}_J$, $\mathfrak{z}(\mathfrak{r}_W) = \mathfrak{h}_J$, $\tilde{\mathfrak{s}}_W = \mathfrak{s}_J$, $\mathfrak{p}_J = \mathfrak{r}_W \oplus \mathfrak{u}_J$. In particular R_W is a Levi subgroup of P_W .
- b) τ is trivial on $\mathfrak{u}_J$.
- c) The representations $\tilde{\tau}_W$ and τ_W are irreducible.
- d) If $J_1, \dots, J_r$ are the connected components of J , then S_{J_i} are the simple factors of $\tilde{S}_W$.
- e) $\mathfrak{h}^{J_i}$ is a Cartan subalgebra of $\text{Lie } S_i = \mathfrak{s}_{J_i}$ and $\mathfrak{h}^J = \bigoplus_{i=1}^r \mathfrak{h}^{J_i}$ is a Cartan subalgebra of $\tilde{\mathfrak{s}}_W$.

Assume by reordering that S_{J_i} acts nontrivially on W if and only if $i \leq q$. Set $I := J_1 \sqcup \cdots \sqcup J_q$, $I' := J_{q+1} \sqcup \cdots \sqcup J_r$. Denote by μ_τ the highest weight of τ with respect to $(\mathfrak{h}, \Pi)$. Then

- f) $\mathfrak{s}_W = \mathfrak{s}_I$, $\mathfrak{s}'_W = \mathfrak{s}_{I'}$, $\mathfrak{h}^J$ is Cartan subalgebra of $\mathfrak{s}_J$, $\mathfrak{h}^I$ is Cartan subalgebra of $\mathfrak{s}_W$ and $\mathfrak{h}^{I'}$ is Cartan subalgebra of $\mathfrak{s}'_W$.
- g) $\Pi' := \{\alpha|_{\mathfrak{h}^J} : \alpha \in J\}$ is a system of simple roots for $(\mathfrak{s}_J, \mathfrak{h}^J)$ and $\Pi'' := \{\alpha|_{\mathfrak{h}^I} : \alpha \in I\}$ is a system of simple roots for $(\mathfrak{s}_W, \mathfrak{h}^I)$.
- h) $\mathfrak{h}^J = \mathfrak{h}^I \oplus \mathfrak{h}^{I'}$, $\mathfrak{h} = \mathfrak{h}_J \oplus \mathfrak{h}^I \oplus \mathfrak{h}^{I'}$, $\mathfrak{h}_I = \mathfrak{h}_J \oplus \mathfrak{h}^{I'}$, $\mathfrak{h}_{I'} = \mathfrak{h}_J \oplus \mathfrak{h}^I$.
- i) If v_τ is a highest weight vector of τ with respect to $(\mathfrak{h}, \Pi)$, then $v_\tau \in W$ and v_τ is also a highest weight vector of $\tilde{\tau}_W$ with respect to $(\mathfrak{h}^J, \Pi')$ and of τ_W with respect to $(\mathfrak{h}^I, \Pi'')$.
- j) If $x_0 = [v_\tau]$ then $\tilde{S}_W \cdot x_0 = S_W \cdot x_0 \subset M_W$; in particular $M_W \neq \emptyset$;
- k) I is μ_τ -connected and J is its μ_τ -saturation.
- l) $W = V_I$.

Proof. (a) The first statement follows by writing elements of $\mathfrak{g}$ in a basis adapted to the compact form as e.g. in [22, pp. 352ff]). The rest follows immediately. (b) Since the representation of P_W on W is irreducible and $[\mathfrak{p}_W, \mathfrak{u}_J] \subset \mathfrak{u}_J$, it follows from Engel theorem that $U_J = \exp \mathfrak{u}_J$ acts trivially and that the representation of R_W on W is irreducible. (c) By Schur lemma the elements of $\exp \mathfrak{h}_J$ act as scalars and the representation $\tilde{\tau}_W$ is irreducible. Since $\tau_W(S_W) = \tilde{\tau}_W(\tilde{S}_W)$ by the definition of S_W , the representation τ_W is irreducible as well. (d)-(g) follow immediately from the definitions of $\mathfrak{s}_J$, $\mathfrak{s}_W$ and $\mathfrak{s}'_W$, see (11). (h) By construction $J = I \sqcup I'$ and $I \perp I'$, so $\mathfrak{h}^J = \mathfrak{h}^I \oplus \mathfrak{h}^{I'}$. Since $\mathfrak{h} = \mathfrak{h}_J \oplus \mathfrak{h}^J$, the second statement follows. Next observe that $\mathfrak{h}_J \subset \mathfrak{h}_I$, since $I \subset J$, and $I' \perp I$, so $\mathfrak{h}^{I'} \subset \mathfrak{h}_I$. Since $\mathfrak{h}_J \perp \mathfrak{h}^{I'}$, $\dim \mathfrak{h}_J \oplus \mathfrak{h}^{I'} = \dim \mathfrak{h} - |J| + |I'| = \dim \mathfrak{h}_I$. This proves the third statement and the same argument yields the fourth. (i) In passing from τ to $\tilde{\tau}_W$ we restrict

both the group and the space. Therefore some care is needed since a priori we don't know that v_τ belongs to the smaller space, i.e. W . So we start by fixing a highest weight vector $w \in W$ of $\tilde{\tau}_W$ with respect to $\mathfrak{h}^J$. Since any element of $\mathfrak{h}_J$ acts on W by scalar multiplication, w is an eigenvector of $\mathfrak{h} = \mathfrak{h}_J \oplus \mathfrak{h}^J$. To check that w is a highest weight of τ with respect to $\mathfrak{h}$ and Π we need to show that $\tau(\mathfrak{g}_\alpha)w = 0$ for any $\alpha \in \Delta_+$. If $\alpha \in \Delta_+$, then $\mathfrak{g}_\alpha \subset \mathfrak{p}_W$ so either $\mathfrak{g}_\alpha \subset \mathfrak{u}_J$ or $\mathfrak{g}_\alpha \subset \mathfrak{s}_W = \mathfrak{s}_J$. In the first case $\tau(\mathfrak{g}_\alpha)w = 0$ since τ is trivial on $\mathfrak{u}_J$ by (b). In the second case $\tau(\mathfrak{g}_\alpha)w = 0$ since w is the highest weight vector of $\tilde{\tau}_W$ and $\alpha \in \Delta_{J,+}$. This shows that $\tau(\mathfrak{g}_\alpha)w = 0$ for any $\alpha \in \Delta_+$ and since τ is irreducible we get that w is also a highest weight of τ , so we denote it by v_τ . In passing from $\tilde{\tau}_W$ to τ_W we only restrict the group (not the representation space!) so it is immediate that v_τ is also a highest weight vector of τ_W with respect to $\mathfrak{h}^I = \mathfrak{h}^J \cap \mathfrak{s}_I$ and Π'' . (j) By the definition of S_W we have $\tilde{S}_W \cdot x_0 = S_W \cdot x_0 \subset \mathbb{P}(W)$ and of course $\tilde{S}_W \cdot x_0 \subset G \cdot x_0$. By Borel-Weil theorem $G \cdot x_0 = M$ since v_τ is a highest weight vector. (k) We need to show that μ_τ is orthogonal to J_i iff $i > q$. By [16, p.197] there are irreducible representations $\sigma_i : S_i \rightarrow \mathrm{GL}(V_i)$ for $i = 1, \dots, r$, such that $W = V_1 \otimes \dots \otimes V_r$ and $\tilde{\tau}_W = \sigma_1 \hat{\otimes} \dots \hat{\otimes} \sigma_r$. The factor S_i acts trivially on W iff σ_i is the trivial representation, which is equivalent to $\mu_\tau|_{\mathfrak{h}^{J_i}} = 0$ i.e. to μ_τ being orthogonal to J_i . So indeed μ_τ is orthogonal to J_i iff $i > q$. (l) We just proved that $W = V_1 \otimes \dots \otimes V_q$, $\tau_W = \sigma_1 \hat{\otimes} \dots \hat{\otimes} \sigma_q$ and $S_I = S_W$. So both W and V_I are irreducible S_I -submodules of V . Since v_τ belongs to both, they must coincide. $\square$

In Theorem 41 we will show that equality holds in (j).

Corollary 32. *If $W \subset V$ is τ -connected, the linear span of M_W is $\mathbb{P}(W)$. If $W_1, W_2 \subset V$ are τ -connected subspaces, then $W_1 \subset W_2$ if and only if $M_{W_1} \subset M_{W_2}$.*

Proof. M_W is a nondegenerate subvariety of $\mathbb{P}(W)$ since it contains the closed orbit $S_W \cdot x_0$ of the irreducible representation τ_W . The result follows. $\square$

Proposition 33. *Let $(\mathfrak{h}, \Pi)$ be a root datum and let μ_τ be the highest weight of τ . Fix a μ_τ -connected subset $I \subset \Pi$ and $g \in G$. Then*

- (a) $W = \tau(g)V_I$ is a τ -connected subspace.
- (b) There is some $a \in K$ such that $W = \tau(a)V_I$.
- (c) $P_W = gP_Jg^{-1}$, $S_W = gS_Ig^{-1}$ and $\tilde{S}_W = gS_Jg^{-1}$, where J is the μ_τ -saturation of I .

Proof. It follows from Lemmata 20 and 22 and from the analysis in the proof of Proposition 31 that V_I is a τ -connected subspace and that $S_{V_I} = S_I$, $P_{V_I} = P_J$. If $W = \tau(g)V_I$ then $P_W = gP_Jg^{-1}$ so in particular P_W is parabolic. Since $G = KP_J$, $g = ap$ for some $a \in K$, $p \in P_J$. Therefore $W = \tau(a)V_I$ and $P_W = aP_Ja^{-1}$ acts irreducibly on W , so W is τ -connected. This proves (a) and (b). (c) follows immediately. $\square$

Corollary 34. *Let $(\mathfrak{h}, \Pi)$ be a root datum and let μ_τ be the highest weight of τ . A subspace $W \subset V$ is τ -connected if and only if $W = \tau(a)V_I$ for some $a \in K$ and for some μ_τ -connected subset $I \subset \Pi$. In this case $P_W = aP_Ja^{-1}$ and $S_W = aS_Ia^{-1}$, where J denotes the μ_τ -saturation of I .*

We can now reformulate Satake's analysis of the boundary of $\overline{X}_\tau^S$. If $W \subset V$ is a τ -connected subspace, the data $S_W, K \cap S_W, \tau_W, \langle \cdot, \cdot \rangle$ are again of the type

described at p. 4. So we can set $X_W := S_W/K \cap S_W$ and there is an embedding analogous to (16)

$$(35) \quad i_{\tau_W} : X_W \rightarrow \mathcal{P}(W) \quad i_{\tau_W}(gK_W) = [\tau_W(g)\tau_W(g)^*].$$

Define

$$i_W := \psi \circ i_{\tau_W} : X_W \rightarrow \overline{\mathcal{P}(V)}$$

where ψ is as in (23). Theorem 24 and Lemma 25 can be rephrased in the following way.

Theorem 36. *The boundary components of $\overline{X}_\tau^S$ are exactly the subsets of $\overline{X}_\tau^S$ of the form $i_W(X_W)$ for some τ -connected subspace $W \subsetneq V$, while $X = i_V(X_V)$. Moreover*

$$(37) \quad \overline{X}_\tau^S = \bigsqcup_{W \text{ } \tau\text{-connected}} i_W(X_W).$$

2.4. Projections and rational maps. In this § we will study the projection $\mathbb{P}(V) \dashrightarrow \mathbb{P}(W)$ induced by the decomposition $V = W \oplus W^\perp$, where W is a τ -connected subspace. Next we will interpret elements of $\overline{X}_\tau^S$ as rational self-maps of M .

For any τ -connected subspace W , denote by $\pi_W : V \rightarrow W$ the orthogonal projection and by $\tilde{\pi}_W$ its projectivization:

$$(38) \quad \tilde{\pi}_W : \mathbb{P}(V) - \mathbb{P}(W^\perp) \rightarrow \mathbb{P}(W) \quad \tilde{\pi}_W([v]) = [\pi_W(v)].$$

Both π_W and $\tilde{\pi}_W$ are S_W -equivariant, since the splitting $V = W \oplus W^\perp$ is preserved by K_W .

Let $(\mathfrak{h}, \Pi)$, μ_τ , v_τ , x_0 , I and J be as in Proposition 31. Let P be the stabilizer of x_0 and let $E \subset \Pi$ be such that $P = P_E$. Set

$$\mathfrak{u}^- := \bigoplus_{\alpha \in \Delta_- - \Delta_E} \mathfrak{g}_\alpha \quad \Delta_{I,-} := \Delta_I \cap \Delta_- \quad \mathfrak{u}^{I,-} := \bigoplus_{\alpha \in \Delta_{I,-} - \Delta_E} \mathfrak{g}_\alpha.$$

By construction $\mathfrak{u}^{I,-} \subset \mathfrak{u}^-$. Let $\mu : \mathfrak{u}^- \rightarrow \mathfrak{u}^{I,-}$ be the projection according to the root space decomposition: if $X = \sum X_\beta \in \mathfrak{u}^-$ with $X_\beta \in \mathfrak{g}_\beta$, then

$$\mu(X) := \sum_{\beta \in \Delta_{I,-} - \Delta_E} X_\beta.$$

Lemma 39. *If $X \in \mathfrak{u}^-$ and $k \in \mathbb{N}$, then $\pi_W(\tau(X)^k \cdot v_\tau) = \tau(\mu(X))^k \cdot v_\tau$.*

Proof. Since $v_\tau \in V_I = W$, $\pi_W(v_\tau) = v_\tau$ so the statement is true for $k = 0$. If $k > 0$ write $X = \sum_{\beta \in \Delta_- - \Delta_E} X_\beta$ with $X_\beta \in \mathfrak{g}_\beta$. Then

$$\tau(X)^k \cdot v_\tau = \sum_{\beta_1, \dots, \beta_k \in \Delta_- - \Delta_E} \tau(X_{\beta_1}) \cdots \tau(X_{\beta_k}) \cdot v_\tau$$

Since $\tau(X_{\beta_1}) \cdots \tau(X_{\beta_k}) \cdot v_\tau \in V_{\mu_\tau + \beta_1 + \dots + \beta_k}$,

$$\pi_W(\tau(X_{\beta_1}) \cdots \tau(X_{\beta_k}) \cdot v_\tau) = \begin{cases} \tau(X_{\beta_1}) \cdots \tau(X_{\beta_k}) \cdot v_\tau & \text{if } \text{supp}(\beta_1 + \dots + \beta_k) \subset I \\ 0 & \text{otherwise.} \end{cases}$$

All the β_i 's are negative roots, so if one of them has nonzero component in the direction of some $\alpha \in I$, this component survives in the sum. Hence $\text{supp}(\beta_1 + \dots + \beta_k) \subset I$ iff $\beta_i \in \Delta_I$ for $i = 1, \dots, k$. Therefore

$$\pi_W(\tau(X)^k \cdot v_\tau) = \sum_{\beta_1, \dots, \beta_k \in \Delta_{I, -\Delta_E}} \tau(X_{\beta_1}) \cdots \tau(X_{\beta_k}) \cdot v_\tau.$$

On the other hand

$$\begin{aligned} \mu(X) &= \sum_{\beta \in \Delta_{I, -\Delta_E}} X_\beta \\ \tau(\mu(X))^k \cdot v_\tau &= \sum_{\beta_1, \dots, \beta_k \in \Delta_{I, -\Delta_E}} \tau(X_{\beta_1}) \cdots \tau(X_{\beta_k}) \cdot v_\tau = \pi_W(\tau(X)^k \cdot v_\tau). \end{aligned}$$

This proves the lemma. $\square$

Set

$$U^- := \exp \mathfrak{u}^- \quad U^{I,-} := \exp \mathfrak{u}^{I,-}.$$

Lemma 40. $\mathfrak{u}^-$ and $\mathfrak{u}^{I,-}$ are nilpotent subalgebras of $\mathfrak{g}$. U^- and $U^{I,-}$ are closed connected algebraic subgroups of G and $S_W = S_I$ respectively. $U^- \cap P = \{e\}$. The maps $\exp : \mathfrak{u}^- \rightarrow U^-$ and $\exp : \mathfrak{u}^{I,-} \rightarrow U^{I,-}$ are diffeomorphisms. The map $a \mapsto \tau(a) \cdot x_0$ is a dominant map of U^- to M and its restriction to $U^{I,-}$ is a dominant map of $U^{I,-}$ to $S_W \cdot x_0$.

For a proof see e.g. [29, p. 51] or [1, p. 68].

Theorem 41. For any τ -connected subspace W , $\tilde{\pi}_W(M - \mathbb{P}(W^\perp)) = S_W \cdot x_0 = M_W$.

Proof. If $X \in \mathfrak{u}^-$ then

$$\tau(\exp X) = \sum_{k=0}^{\infty} \frac{\tau(X)^k}{k!} \quad \tau(\exp \mu(X)) = \sum_{k=0}^{\infty} \frac{\tau(\mu(X))^k}{k!}$$

and the sums are finite since $\tau(X)$ and $\tau(\mu(X))$ are nilpotent. Using Lemma 39 we get

$$\begin{aligned} \pi_W(\tau(\exp X) \cdot v_\tau) &= \sum_{k=0}^{\infty} \frac{1}{k!} \pi_W(\tau(X)^k \cdot v_\tau) = \sum_{k=0}^{\infty} \frac{1}{k!} \tau(\mu(X))^k \cdot v_\tau = \\ &= \exp \tau(\mu(X)) \cdot v_\tau = \tau(\exp \mu(X)) \cdot v_\tau. \end{aligned}$$

This proves that

$$(42) \quad \pi_W(U^- \cdot v_\tau) \subset U^{I,-} \cdot v_\tau.$$

Since $v_\tau \neq 0$, $0 \notin \pi_W(U^- \cdot v_\tau)$, i.e. $U^- \cdot v_\tau \cap W^\perp = \emptyset$. Therefore $U^- \cdot x_0 \subset \mathbb{P}(V) - \mathbb{P}(W^\perp)$. By Lemma 40 $U^- \cdot x_0$ is dense in $G \cdot x_0$. Hence it follows from (42) that

$$(43) \quad \tilde{\pi}_W(M - \mathbb{P}(W^\perp)) \subset S_W \cdot x_0.$$

On the other hand S_W preserves $\mathbb{P}(W)$, so $S_W \cdot v_\tau \cap W^\perp = \emptyset$ and

$$S_W \cdot x_0 = \tilde{\pi}_W(S_W \cdot x_0) \subset \tilde{\pi}_W(M - \mathbb{P}(W^\perp)).$$

This proves that $\tilde{\pi}_W(M - \mathbb{P}(W^\perp)) = S_W \cdot x_0$. If $g \in S_W$, then $g \cdot x_0 \in G \cdot x_0 = M$ and $g \cdot x_0 \in \mathbb{P}(W)$ since S_W preserves $W = V_I$. Hence $S_W \cdot x_0 \subset M \cap \mathbb{P}(W) = M_W$.

Conversely, if $x \in M_W$, then $x \notin \mathbb{P}(W^\perp)$ and $\tilde{\pi}_W(x) = x$. But $x \in M$, so $\tilde{\pi}_W(x) \in S_W \cdot x_0$ by (43). Therefore $M_W \subset S_W \cdot x_0$. This proves that $S_W \cdot x_0 = M_W$. $\square$

We can now give the interpretation in terms of rational maps. For technical reasons that will become clear later (Theorem 85) we prefer to take a square root. Recall the following elementary fact.

Lemma 44. *Let V be a Hermitian vector space and set $\mathcal{S}(V) = \{A \in \mathcal{H}(V) : A \geq 0\}$. If $A \in \mathcal{S}(V)$ there is a unique $B \in \mathcal{S}(V)$ such that $B^2 = A$. Set $\sqrt{A} := B$. Then $\sqrt{\cdot}$ is a homeomorphism of $\mathcal{S}(V)$ onto itself.*

Proof. It is enough to prove that $q : \mathcal{S}(V) \rightarrow \mathcal{S}(V)$, $q(A) = A^2$ is a homeomorphism. q is of course continuous and using the spectral theorem one easily proves that it is bijective. It is enough to show that q is proper. Let $\{A_n\}$ be a sequence in $\mathcal{S}(V)$ such that $q(A_n) = A_n^2 \rightarrow B$. Then $\text{tr } A_n^* A_n = \text{tr } A_n^2 \rightarrow \text{tr } B$. Therefore $\text{tr } A_n^* A_n$ is bounded and A_n admits a subsequence that converges to some A , which necessarily lies in $\mathcal{S}(V)$ since $\mathcal{S}(V)$ is closed in $\mathcal{H}(V)$. $\square$

Let G, K, τ and $\langle \cdot, \cdot \rangle$ be as at p. 4. Then

$$\mathfrak{P} := \{g \in G : \theta(g) = g^{-1}\}$$

is a submanifold of G , $\exp : i\mathfrak{k} \rightarrow \mathfrak{P}$ is a diffeomorphism with inverse $\log : \mathfrak{P} \rightarrow i\mathfrak{k}$, the map

$$g \rightarrow \sqrt{g} := \exp\left(\frac{\log g}{2}\right)$$

is a diffeomorphism of $\mathfrak{P}$ onto itself and the map $\mathfrak{P} \rightarrow X$, $g \mapsto gK$ is a diffeomorphism. Set

$$(45) \quad \rho : G \rightarrow \mathfrak{P} \quad \rho(g) := \sqrt{g\theta(g^{-1})}.$$

Then $a = \rho(g)^{-1}g \in K$ and $g = \rho(g) \cdot a$ is the polar decomposition of g .

Definition 46. *If $p = [A] \in \overline{X}_\tau^S$, let $R_p : \mathbb{P}(V) \dashrightarrow \mathbb{P}(V)$ be the rational map induced by the endomorphism $\sqrt{A} \in \mathcal{S}(V)$, i.e. R_p is defined outside of $\mathbb{P}(\ker A)$ and for $x = [v] \notin \mathbb{P}(\ker A)$, $R_p(x) = [\sqrt{A}v]$. Set*

$$(47) \quad \check{p} := R_p|_M : M \dashrightarrow M.$$

For this to make sense we need to show that M intersects the domain of definition of R_p and that $R_p(M) \subset M$.

Lemma 48. *Fix $g \in S_W$ and set $p = i_W(gK_W) \in \overline{X}_\tau^S$. The following hold.*

- a) *The indeterminacy locus of R_p is $\mathbb{P}(W^\perp)$.*
- b) *For $x = [v] \in \mathbb{P}(V) - \mathbb{P}(W^\perp)$*

$$R_p(x) = \left[\sqrt{\tau(g)\tau(g)^*} \pi_W(v) \right].$$

- c) *$M_W \subset M - \mathbb{P}(W^\perp)$.*
- d) *If $x \in M_W$, then $R_p(x) = \rho(g) \cdot x$.*
- e) *$R_p(M - \mathbb{P}(W^\perp)) = M_W$.*
- f) *$\check{p}$ is a map as in (47) and $\check{p} = L_{\rho(g)} \circ \tilde{\pi}_W$, where $L_{\rho(g)}$ denotes the automorphism of M_W induced by $\rho(g)$.*
- g) *If $p \in \overline{X}_\tau^S$, then $p \in i_W(X_W)$ iff $\text{Im } \check{p} = M_W$.*

Proof. (a) Set $A = \tau_W(g)\tau_W(g)^* \oplus 0$. Then $p = [A]$ and R_p is the rational map defined by the endomorphism $\sqrt{A} \in \text{End } V$. Therefore the indeterminacy locus is $\mathbb{P}(\ker A) = \mathbb{P}(W^\perp)$. (b) and (c) are obvious. (d) Set $g_1 = \rho(g)$, $a = g_1^{-1}g$. Then $g\theta(g^{-1}) = g_1^2$, $A = \tau_W(g)\tau_W(g)^* = \tau_W(g_1)^2$, so $\sqrt{A} = \tau_W(g_1)$. If $x = [v] \in M_W$, then $R_p(x) = [\sqrt{A}\pi_W(v)] = [\tau_W(g_1)v] = g_1 \cdot x$. (e) By what we just proved and Theorem 41 $R_p(M - \mathbb{P}(W^\perp)) = L_{g_1}\tilde{\pi}_W(M - \mathbb{P}(W^\perp)) = L_{g_1}(M_W) = M_W$. (f) By (c) $M \not\subset \mathbb{P}(W^\perp)$ so $\check{p} = R_p|_M$ is a well-defined rational map and by (e) its image is contained in M . The rest is just a restatement of (e). (g) This follows from (e) and Corollary 32. $\square$

2.5. τ -connected subspaces and the moment map. In this § we study τ -connected subspaces from the point of view of the coadjoint orbit $\mathcal{O} = \Phi(M)$. We will show that to each τ -connected subspace W there corresponds a set of affine inequalities that are satisfied by the points of the coadjoint orbit. Moreover the points where the equalities hold are exactly the points of $\Phi(M_W)$.

Definition 49. *Let W be a τ -connected subspace and let $(\mathfrak{h}, \Pi)$ be a root datum. We say that $(\mathfrak{h}, \Pi)$ is a W -datum if $\theta(\mathfrak{h}) = \mathfrak{h}$ and $\mathfrak{b}_+(\mathfrak{h}, \Pi) \subset \mathfrak{p}_W$. The set of W -data will be denoted by $\mathcal{D}(W)$.*

By Corollary 28 any τ -connected subspace admits a W -datum, which in general is not unique. If a W -datum is fixed, then the following objects are well-defined: a subset $J \subset \Pi$ such that $\mathfrak{p}_W = \mathfrak{p}_J$, a subset $I \subset J$ such that $W = V_I$, a highest weight $\mu_\tau \in \mathfrak{h}^*$ and a line $x_0 = [v_\tau] \in M$ of highest weight vectors. When we use these symbols we understand that a W -datum has been chosen and that they refer to that particular choice. By (i) of Prop. 31 $x_0 \in M_W$. Any point $x_0 \in M_W$ is obtained in this way from a W -datum, but in general in many ways. In other words, the map $(\mathfrak{h}, \Pi) \mapsto x_0$ from the set of W -data to M_W is surjective, but in general non-injective.

Lemma 50. *Let $W \subset V$ be a τ -connected subspace and let $(\mathfrak{h}, \Pi)$ be a W -datum. Then the nonzero weights of the adjoint representation of $\mathfrak{z}_{\mathfrak{g}}(\mathfrak{r}_W)$ on $\mathfrak{p}_W$ coincide with the nonzero restrictions to $\mathfrak{z}_{\mathfrak{g}}(\mathfrak{r}_W)$ of elements of $\Delta_+(\mathfrak{h}, \Pi)$. In particular the set of these restrictions does not depend on the choice of $(\mathfrak{h}, \Pi)$, but only on W .*

Proof. By (a) of Prop. 31 $\mathfrak{z}_{\mathfrak{g}}(\mathfrak{r}_W) = \mathfrak{h}_J$. The decomposition

$$\mathfrak{p}_W = \mathfrak{r}_W \oplus \bigoplus_{\alpha \in \Delta_+ - \Delta_J} \mathfrak{g}_\alpha$$

is clearly $\text{ad } \mathfrak{h}_J$ -invariant. For $\lambda \in \mathfrak{h}_J^*$ denote by U_λ the corresponding weight space of $\text{ad} : \mathfrak{h}_J \rightarrow \mathfrak{gl}(\mathfrak{p}_W)$. If $\alpha \in \Delta$, then $\alpha|_{\mathfrak{h}_J} = 0$ if and only if $\alpha \in \Delta_J$. So $U_0 = \mathfrak{r}_W$ and for λ a nonzero weight $U_\lambda = \bigoplus_{\alpha \in \Delta_+ : \alpha|_{\mathfrak{h}_J} = \lambda} \mathfrak{g}_\alpha$. $\square$

Let W be a τ -connected subspace and let $(\mathfrak{h}, \Pi)$ be a W -datum. Let $\mathfrak{k}'_W$ be as in (30). Set

$$\mathfrak{c}_W := \mathfrak{z}_{\mathfrak{t}}(\mathfrak{r}_W) \oplus \mathfrak{k}'_W \quad \mathfrak{c}_{W, \mathfrak{h}, \Pi} := \mathfrak{h}_I \cap \mathfrak{k}.$$

By definition $\mathfrak{c}_W$ depends only on W , while $\mathfrak{c}_{W, \mathfrak{h}, \Pi}$ does in general depend also on $(\mathfrak{h}, \Pi)$. Since $\mathfrak{c}_W = (\mathfrak{h}_J \cap \mathfrak{k}) \oplus \mathfrak{k}_{I'}$ and $\mathfrak{h}_I = \mathfrak{h}_J \oplus \mathfrak{h}^{I'}$ by Prop. 31 (h), it follows that $\mathfrak{c}_{W, \mathfrak{h}, \Pi} \subset \mathfrak{c}_W$ for any W -datum $(\mathfrak{h}, \Pi)$.

Lemma 51. *If $x, x' \in M_W$, then $\Phi(x)|_{\mathfrak{c}_W} = \Phi(x')|_{\mathfrak{c}_W}$.*

Proof. $x' = a \cdot x$ for some $a \in K_W$. $[\mathfrak{k}_W, \mathfrak{c}_W] = 0$ and K_W is connected. So if $v \in \mathfrak{c}_W$, then $\text{Ad}(a^{-1})v = v$ and $\langle \Phi(x'), v \rangle = \langle \Phi(x), \text{Ad}(a^{-1})v \rangle = \langle \Phi(x), v \rangle$. $\square$

Pick $x \in M_W$ and set

$$E_W := \{\lambda \in \mathfrak{k}^* : \lambda(v) = \langle \Phi(x), v \rangle \text{ for all } v \in \mathfrak{c}_W\}.$$

By the previous lemma E_W does not depend on the choice of $x \in M_W$. Similarly, for a W -datum $(\mathfrak{h}, \Pi)$ define

$$(52) \quad E_{W, \mathfrak{h}, \Pi} := \{\lambda \in \mathfrak{k}^* : \lambda(v) = \langle \Phi(x_0), v \rangle \text{ for all } v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}\}.$$

(As usual x_0 is the line thorough the highest weight vector determined by $(\mathfrak{h}, \Pi)$.)

Lemma 53.

$$E_W = \bigcap_{(\mathfrak{h}, \Pi) \in \mathcal{D}(W)} E_{W, \mathfrak{h}, \Pi}.$$

Proof. Since $\mathfrak{c}_{W, \mathfrak{h}, \Pi} \subset \mathfrak{c}_W$, $E_W \subset E_{W, \mathfrak{h}, \Pi}$. One inclusion follows. For the other take $\lambda \in \bigcap E_{W, \mathfrak{h}, \Pi}$. If $v \in \mathfrak{c}_W$, then $v = v_1 + v_2 \in \mathfrak{z}_{\mathfrak{k}}(\mathfrak{r}_W) \oplus \mathfrak{k}'_W$. v_2 lies in the Lie algebra $\mathfrak{t}_2$ of a maximal torus of $\mathfrak{k}'_W$. Completing $\mathfrak{z}_{\mathfrak{k}}(\mathfrak{r}_W) \oplus (\mathfrak{t}_2 \otimes \mathbb{C})$ to a Cartan subalgebra of $\mathfrak{g}$ we get a datum $(\mathfrak{h}, \Pi)$ such that $v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}$. Since $\lambda \in E_{W, \mathfrak{h}, \Pi}$ we get $\lambda(v) = \langle \Phi(x_0), v \rangle$. This proves that $\lambda \in E_W$. $\square$

Next set

$$\mathfrak{c}_{W, \mathfrak{h}, \Pi}^- := \{v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi} : i\alpha(v) \leq 0 \text{ for all } \alpha \in \Pi\}.$$

Note that $\mathfrak{c}_{W, \mathfrak{h}, \Pi}^- = \{v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi} : i\alpha(v) \leq 0 \text{ for all } \alpha \in \Pi - I\}$, since $\mathfrak{c}_{W, \mathfrak{h}, \Pi} = \bigcap_{\alpha \in I} \ker i\alpha \cap \mathfrak{k}$. Moreover $\{i\alpha : \alpha \in \Pi\}$ is a basis of $\mathfrak{k}^*$ and $\{i\alpha|_{\mathfrak{c}_{W, \mathfrak{h}, \Pi}} : \alpha \in \Pi - I\}$ is a basis of $\mathfrak{c}_{W, \mathfrak{h}, \Pi}^*$. Therefore $\mathfrak{c}_{W, \mathfrak{h}, \Pi}^-$ is just a (higher dimensional) quadrant, so in particular it is a convex cone with nonempty interior spanned by a finite number of rays. Put

$$(54) \quad \begin{aligned} C_{W, \mathfrak{h}, \Pi} &:= \{\lambda \in \mathfrak{k}^* : \lambda(v) \leq \langle \Phi(x_0), v \rangle \text{ for all } v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}^-\} \\ C_W &:= \bigcap_{(\mathfrak{h}, \Pi) \in \mathcal{D}(W)} C_{W, \mathfrak{h}, \Pi} \quad A_W := \bigcup_{(\mathfrak{h}, \Pi) \in \mathcal{D}(W)} \mathfrak{c}_{W, \mathfrak{h}, \Pi}^- \end{aligned}$$

Lemma 55. *Let x be any point in M_W . Then $C_W = \{\lambda \in \mathfrak{k}^* : \lambda(v) \leq \langle \Phi(x), v \rangle \text{ for all } v \in A_W\}$ and $E_W = \{\lambda \in \mathfrak{k}^* : \lambda(v) = \langle \Phi(x), v \rangle \text{ for all } v \in A_W\}$.*

Proof. The choice of x is irrelevant, by Lemma 51 since $v \in \mathfrak{c}_W$. The first statement follows directly from the definitions of C_W and A_W . Moreover $\mathfrak{c}_{W, \mathfrak{h}, \Pi}^-$ spans $\mathfrak{c}_{W, \mathfrak{h}, \Pi}$, so $E_{W, \mathfrak{h}, \Pi} = \{\lambda \in \mathfrak{k}^* : \lambda(v) = \langle \Phi(x), v \rangle \text{ for all } v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}^-\}$. So the second statement follows from Lemma 53. Put in another way, the linear span of A_W is $\mathfrak{c}_W$. $\square$

For $v \in \mathfrak{k}$ set $h_v := \langle \Phi, v \rangle$. h_v is a component of the moment map and it is the Hamiltonian function of ξ_v , i.e. $-i_{\xi_v}\omega = dh_v$. Let H_{μ_τ} be as in (9).

Lemma 56. *Let $W \subset V$ be τ -connected and let $(\mathfrak{h}, \Pi)$ be a W -datum.*

- a) *If $v \in \mathfrak{c}_W$, then x_0 is a critical point of h_v .*
- b) *If $v \in \mathfrak{c}_W$, $u \in \mathfrak{k}$ and $w = \xi_u(x_0)$, then*

$$D^2 h_v(x_0)(w, w) = iB([u, H_{\mu_\tau}], [u, v]).$$

Proof. Since K is transitive on M any $w \in T_{x_0}M$ can be written as $w = \xi_u(x_0)$ for some $u \in \mathfrak{k}$. Set $\alpha(t) = \exp(tu) \cdot x_0$ and $\gamma(t) = \text{Ad}(\exp(-tu))v$. Then

$$h_v(\alpha(t)) = \langle \Phi(\exp tu \cdot x_0), v \rangle = \langle \Phi(x_0), \text{Ad}(\exp(-tu))v \rangle = \langle \Phi(x_0), \gamma(t) \rangle$$

$$dh_v(x_0)(w) = \left. \frac{d}{dt} \right|_{t=0} h_v(\alpha(t)) = \langle \Phi(x_0), \dot{\gamma}(0) \rangle.$$

Clearly $\dot{\gamma}(0) = -[u, v]$, so using (13)

$$dh_v(x_0)(w) = i\mu_\tau([u, v]^\mathfrak{h}) = B(iH_{\mu_\tau}, [u, v]) = iB([v, H_{\mu_\tau}], u).$$

Since I' is orthogonal to μ_τ , $H_{\mu_\tau} \in \mathfrak{h}_{I'}$ and $\mathfrak{h}_{I'} = \mathfrak{h}_J \oplus \mathfrak{h}^I \subset \mathfrak{h}_J \oplus \mathfrak{k}_I$ by Prop 31 (h). Hence $[\mathfrak{c}_W, H_{\mu_\tau}] = 0$. Thus $dh_v(x_0) = 0$ and (a) is proved. To compute the Hessian consider again the path α . Since x_0 is critical

$$D^2h_v(x_0)(w, w) = \left. \frac{d^2}{dt^2} \right|_{t=0} h_v(\alpha(t)) = \langle \Phi(x_0), \ddot{\gamma}(0) \rangle = \langle \Phi(x_0), [u, [u, v]] \rangle =$$

$$= -iB(H_{\mu_\tau}, [u, [u, v]]) = iB([u, H_{\mu_\tau}], [u, v]).$$

This proves the second statement. $\square$

Theorem 57. *If W is a proper τ -connected subspace, then $\mathcal{O} = \Phi(M) \subset C_W$ and $\Phi^{-1}(E_W) = M_W$.*

Proof. By the definition (54) of C_W , to prove the first part of the theorem we need to show that $\mathcal{O} = \Phi(M) \subset C_{W, \mathfrak{h}, \Pi}$ for any W -datum $(\mathfrak{h}, \Pi)$. Let $(\mathfrak{h}, \Pi)$ be such a datum. It is enough to show that for any $v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}^-$ the maximum of h_v on M is attained at x_0 . We know (see e.g. [22, pp. 352ff]) that it is possible to choose vectors $X_\beta \in \mathfrak{g}_\beta$ for $\beta \in \Delta$ in such a way that as β varies in Δ_+ the vectors

$$(58) \quad A_\beta = \frac{i}{\sqrt{2}}(X_\beta + X_{-\beta}) \quad B_\beta = \frac{1}{\sqrt{2}}(X_\beta - X_{-\beta})$$

describe a $(-B)$ -orthonormal basis of the orthogonal complement of $\mathfrak{k} = \mathfrak{k} \cap \mathfrak{h}$ inside $\mathfrak{k}$. This means that

$$(59) \quad B(A_\beta, B_\gamma) = 0 \quad B(A_\beta, A_\gamma) = B(B_\beta, B_\gamma) = -\delta_{\beta\gamma}.$$

If $H \in \mathfrak{h}$ then $[A_\beta, H] = -i\beta(H)B_\beta$ and $[B_\beta, H] = i\beta(H)A_\beta$. Given $w \in T_{x_0}M$, let $u \in \mathfrak{k}$ be such that $w = \xi_u(x_0)$. Write

$$u = u^\mathfrak{k} + \sum_{\beta \in \Delta_+} (a_\beta A_\beta + b_\beta B_\beta)$$

where $u^\mathfrak{k} \in \mathfrak{k}$. Then for any $H \in \mathfrak{h}$

$$[u, H] = i \sum_{\beta \in \Delta_+} \beta(H) (-a_\beta B_\beta + b_\beta A_\beta).$$

Applying this formula with $H = H_{\mu_\tau}$ and then with $H = v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}^- \subset \mathfrak{h}$, using (59) and applying the computation of the previous lemma, we get

$$D^2h_v(x_0)(w, w) = i \sum_{\beta \in \Delta_+} (a_\beta^2 + b_\beta^2) \beta(H_\tau) \beta(v).$$

If $\beta \in \Delta_+$, then $\beta(H_{\mu_\tau}) \geq 0$ since μ_τ is dominant, while $i\beta(v) \leq 0$ since $v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}^-$. Therefore we get that the Hessian at x_0 is negative semidefinite. But h_v is a component of the moment map $\Phi : M \rightarrow \mathfrak{k}^*$, hence it is a Morse-Bott function with

critical points of even index (this is Frankel theorem, see e.g. [4, Thm. 2.3, p. 109] or [27, p. 186]) and any local maximum point is an absolute maximum point (see e.g. [4, p. 112]). Therefore x_0 is a global maximum of h_v on M . This proves that $\mathcal{O} \subset C_{W, \mathfrak{h}, \Pi}$. Since $(\mathfrak{h}, \Pi)$ is an arbitrary W -datum, we get the first part of the theorem.

Now fix a W -datum $(\mathfrak{h}, \Pi)$ and set

$$F = \{x \in M : h_v(x) = h_v(x_0) \text{ for any } v \in \mathfrak{c}_{W, \mathfrak{h}, \Pi}\}.$$

Recalling definition (52) one immediately recognizes that

$$(60) \quad F = \Phi^{-1}(E_{W, \mathfrak{h}, \Pi}).$$

We start by showing that $F = M_W$. Set for simplicity $\mathfrak{t}_I := \mathfrak{c}_{W, \mathfrak{h}, \Pi}$. The subgroup $T_I = \exp \mathfrak{t}_I$ is a closed torus in K since $\mathfrak{t}_I \otimes \mathbb{C} = \mathfrak{h}_I$ is a Cartan subalgebra of $\mathfrak{s}_I$. The moment map for the action of T_I on M is

$$\Phi_I : M \rightarrow \mathfrak{t}_I^* \quad \Phi_I(x) := \Phi(x)|_{\mathfrak{t}_I}.$$

Set $\lambda = \Phi_I(x_0)$. It is immediate that $F = \Phi_I^{-1}(\lambda)$. By Atiyah theorem [3, Thm. 1.A] every fibre of Φ_I is connected, so F is connected. Let $v_1, \dots, v_r$ be a basis of $\mathfrak{c}_{W, \mathfrak{h}, \Pi}$ such that each v_j lies in $\mathfrak{c}_{W, \mathfrak{h}, \Pi}^-$. Each function h_{v_j} attains its maximum at x_0 , hence the point λ is a vertex of $\Phi_I(M) \subset \mathfrak{t}_I^*$. By the Atiyah-Guillemin-Sternberg convexity theorem F is a connected (symplectic) submanifold. By Lemma 51 we have $\langle \Phi(x), v \rangle = \langle \Phi(x_0), v \rangle$ for any $x \in M_W$ and any $v \in \mathfrak{t}_I$, so $\Phi_I(M_W) = \{\lambda\}$, i.e. $M_W \subset F$. Since M_W and F are compact connected submanifolds, to conclude that $F = M_W$ it is enough to check that $T_{x_0}F = T_{x_0}M_W$. Denote by P the stabilizer of x_0 and by $L = P \cap K$ its stabilizer in K . Since x_0 is the line of highest weight vectors $\mathfrak{b}_+(\mathfrak{h}, \Pi) \subset \mathfrak{p}$, so $P = P_E$ for some subset $E \subset \Pi$. Let $\mathfrak{m}$ denote the orthogonal complement of $\mathfrak{l}$ inside $\mathfrak{k}$. Then there is an isomorphism $\mathfrak{m} \cong T_{x_0}M$ given by $v \mapsto \xi_v(x_0)$. Since F is the connected component of the fixed point set of T_I through x_0 , the tangent space $T_{x_0}F$ is the fixed set of the isotropy representation of T_I on $T_{x_0}M$. Via the isomorphism $\mathfrak{m} \cong T_{x_0}M$ the isotropy representation is identified with the restriction of $\text{Ad } T_I$ on $\mathfrak{m}$. So $T_{x_0}F = \mathfrak{z}(\mathfrak{t}_I) \cap \mathfrak{m}$. We claim that $T_{x_0}M_W = \mathfrak{k}_W \cap \mathfrak{m}$. To check this it is enough to show that

$$(61) \quad \mathfrak{k}_W = \mathfrak{k}_W \cap \mathfrak{l} \oplus \mathfrak{k}_W \cap \mathfrak{m}.$$

This follows from the following fact: if T is the maximal torus corresponding to $\mathfrak{h}$, then $\mathfrak{k}_W$ and $\mathfrak{l}$ are T -invariant subspaces of $\mathfrak{k}$ since $T \subset L$ and $\mathfrak{k}_W = \mathfrak{k}_I$. So also $\mathfrak{m}$ is T -invariant. Therefore both summands on the right in (61) are T -invariant subspaces and $\mathfrak{t} \cap \mathfrak{m} = \{0\}$. Hence (61) follows from the uniqueness of the real root decomposition. We have proved that $T_{x_0}F = \mathfrak{z}(\mathfrak{t}_I) \cap \mathfrak{m}$ and $T_{x_0}M_W = \mathfrak{k}_W \cap \mathfrak{m}$. To show that $T_{x_0}F = T_{x_0}M_W$, we have to check that $\mathfrak{z}(\mathfrak{t}_I) \cap \mathfrak{m} = \mathfrak{k}_W \cap \mathfrak{m}$. Recall (58) and set $\mathfrak{k}_\beta = \mathbb{R}A_\beta + \mathbb{R}B_\beta$ and $\mathfrak{t}^I = \mathfrak{h}^I \cap \mathfrak{k}$. Then

$$\begin{aligned} \mathfrak{k}_W = \mathfrak{k}_I = \mathfrak{t}^I \oplus \bigoplus_{\beta \in \Delta_I \cap \Delta_+} \mathfrak{k}_\beta \\ \mathfrak{l} = \mathfrak{k} \cap \mathfrak{p}_E = \mathfrak{t} \oplus \bigoplus_{\beta \in \Delta_E \cap \Delta_+} \mathfrak{k}_\beta \quad \mathfrak{m} = \bigoplus_{\beta \in \Delta_+ - \Delta_E} \mathfrak{k}_\beta \end{aligned}$$

We claim that

$$\mathfrak{z}(\mathfrak{h}_I) = \mathfrak{h} \oplus \bigoplus_{\beta \in \Delta_I} \mathfrak{g}_\beta.$$

To prove this it is enough to check that if $\beta \in \Delta_+$ then $\beta|_{\mathfrak{h}_I} = 0$ if and only if $\beta \in \Delta_I$. The collection $\{\alpha|_{\mathfrak{h}_I} : \alpha \notin I\}$ is a basis of $\mathfrak{h}_I^*$. If $\beta = \sum_{\alpha \in \Pi} m_\alpha \alpha$ and $\beta|_{\mathfrak{h}_I} = 0$ then

$$\sum_{\alpha \notin I} m_\alpha \alpha|_{\mathfrak{h}_I} = 0$$

so $m_\alpha = 0$ for any $\alpha \notin I$ and $\beta \in \Delta_I$. The opposite implication is trivial, so the claim is proved. From it we get

$$\mathfrak{z}(\mathfrak{t}_I) = \mathfrak{t} \oplus \bigoplus_{\beta \in \Delta_I \cap \Delta_+} \mathfrak{k}_\beta \quad \mathfrak{z}(\mathfrak{t}_I) \cap \mathfrak{m} = \bigoplus_{\beta \in \Delta_I \cap \Delta_+ - \Delta_E} \mathfrak{k}_\beta = \mathfrak{k}_I \cap \mathfrak{m}.$$

Therefore $T_{x_0}M_W = T_{x_0}F$ and $F = M_W$. By (60) this means that $M_W = \Phi^{-1}(E_{W, \mathfrak{h}, \Pi})$. Using Lemma 53 we get

$$M_W = \Phi^{-1}\left(\bigcap_{W, \mathfrak{h}, \Pi} E_{W, \mathfrak{h}, \Pi}\right) = \Phi^{-1}(E_W).$$

This concludes the proof of the theorem. $\square$

We close this section by relating the moment map of M_W to that of M . Let $W \subset V$ be τ -connected. Denote by $\Phi^W : M_W \rightarrow \mathfrak{k}_W^*$ the moment map of the smaller flag manifold (M_W, ω) . Fix a W -datum $(\mathfrak{h}, \Pi)$ and write $H_{\mu_\tau} = Y_W + Z_W$ where $Y_W \in i\mathfrak{k}_W$ and $Z_W \in (i\mathfrak{k}_W)^\perp$.

Lemma 62. $Z_W \in \mathfrak{h}_I$.

Proof. Set $\mathfrak{t} = \mathfrak{h} \cap \mathfrak{k}$, $\mathfrak{t}^I = \mathfrak{h}^I \cap \mathfrak{k}$, $\mathfrak{t}_I = \mathfrak{h}_I \cap \mathfrak{k}$. The two spaces $i\mathfrak{t}_I$, $i\mathfrak{t}^I$ are orthogonal to each other and $H_{\mu_\tau} \in i\mathfrak{t} = i\mathfrak{t}^I \oplus i\mathfrak{t}_I$. Moreover $i\mathfrak{t}^I \subset i\mathfrak{k}_W$ and $i\mathfrak{t}_I \subset (i\mathfrak{k}_W)^\perp$. Therefore $Z_W \in i\mathfrak{t}_I$. $\square$

Pick $x \in M_W$ and define a map $\Theta_W : \mathfrak{k}_W^* \rightarrow \mathfrak{k}^*$ by the following recipe: if $\lambda \in \mathfrak{k}_W^*$ and $v \in \mathfrak{k}$, write $v = v_1 + v_2$ where $v_1 \in \mathfrak{k}_W$ and $v_2 \in \mathfrak{k}_W^\perp$ and set

$$(63) \quad \Theta_W(\lambda)(v) := \lambda(v_1) + \langle \Phi(x), v_2 \rangle.$$

Θ_W is an injective affine map.

Lemma 64. a) The map Θ_W does not depend on the choice of $x \in M_W$.

b) If $x \in M_W$, then $\Phi(x) = \Theta_W(\Phi^W(x))$.

Proof. Fix $x \in M_W$. To prove (a) it is enough to show that $\langle \Phi(x), v \rangle = \langle \Phi(x_0), v \rangle$ for any $v \in \mathfrak{k}_W^\perp$. There exists $a \in K_W$ such that $x = a \cdot x_0$ and $\langle \Phi(x), v \rangle = \langle \Phi(x_0), \text{Ad}(a^{-1})v \rangle$. So we need to show that $\langle \Phi(x_0), \text{Ad}(a^{-1})v - v \rangle = 0$ for any $a \in K_W$ and any $v \in \mathfrak{k}_W^\perp$. Since K_W is connected it is enough to prove that $\Phi(x_0)|_{[\mathfrak{k}_W, \mathfrak{k}_W^\perp]} = 0$. Note that $\text{Ad } K_W$ preserves the decomposition $\mathfrak{k} = \mathfrak{k}_W \oplus \mathfrak{k}_W^\perp$. By differentiating we infer that $[\mathfrak{k}_W, \mathfrak{k}_W^\perp] \subset \mathfrak{k}_W^\perp$. So if $v \in \mathfrak{k}_W^\perp$ and $u \in \mathfrak{k}_W$, then $[u, v] \in \mathfrak{k}_W^\perp$ and $[u, Z_W] = 0$ by Lemma 62. Therefore using (13)

$$\langle \Phi(x_0), [u, v] \rangle = -iB(H_{\mu_\tau}, [u, v]) = -iB(Z_W, [u, v]) = iB([u, Z_W], v) = 0.$$

This proves (a). (b) is simply a restatement of the fact that for any $x \in M_W$, $\Phi^W(x) = \Phi(x)|_{\mathfrak{k}_W}$. $\square$

3. THE BOURGUIGNON-LI-YAU MAP

3.1. Definition and boundary behaviour. In this section we introduce the Bourguignon-Li-Yau map in the most general case and we investigate its boundary behaviour in the case of a flag manifold. Let K be a connected compact Lie group (not necessarily semisimple) and let $G = K^{\mathbb{C}}$ be its complexification. Set $X = G/K$, $\mathfrak{k} = \text{Lie } K$, $\mathfrak{g} = \text{Lie } G$ and denote by θ the Cartan involution. Let (M, J, g, ω) be a compact Kähler manifold and assume that G acts holomorphically and almost effectively on M (i.e. only finitely many elements of G act trivially on M .) Assume also that the action of K is Hamiltonian with moment map $\Phi : M \rightarrow \mathfrak{k}^*$.

Definition 65. *Given a probability measure γ on M the Bourguignon-Li-Yau map $\Psi_\gamma : X \rightarrow \mathfrak{k}^*$ is defined by*

$$(66) \quad \Psi_\gamma(gK) = \int_M \Phi(\rho(g) \cdot x) d\gamma(x)$$

where $\rho(g) = \sqrt{g\theta(g^{-1})}$ as in (45).

Recall the following notation: if $F : M \rightarrow N$ is a measurable map between measurable spaces and μ is a measure on M , then $F_\# \mu$ denotes the push-forward or image measure. This means that for a measurable subset $E \subset N$ we have $F_\# \mu(E) := \mu(F^{-1}(E))$. For any measurable function φ on N the following change of variable formula holds true

$$(67) \quad \int_N \varphi(y) d(F_\# \mu)(y) = \int_M \varphi(F(x)) d\mu(x)$$

(see e.g. [19, p.163], [23, p.221]). In particular, if M and N are differentiable manifolds, F is a diffeomorphism and μ is the measure associated to a smooth top dimensional form η , then $F_\# \mu$ is the measure associated to the form $(F^{-1})^* \eta$.

Lemma 68. $\Psi_\gamma(X)$ is contained in the convex envelope of $\Phi(M)$ inside $\mathfrak{k}^*$.

Proof. Let $F : M \rightarrow M$ denote multiplication by $\rho(g)$. By formula (67)

$$\Psi_\gamma(gK) = \int_M \Phi(F(x)) d\gamma(x) = \int_M \Phi(y) d(F_\# \gamma)(y) = \int_{\Phi(M)} z d((\Phi \circ F)_\# \gamma)(z).$$

$(\Phi \circ F)_\# \gamma$ is a probability measure supported on $\Phi(M)$ and $\Psi_\gamma(gK)$ is its barycenter, which of course lies in the convex envelope of $\Phi(M)$. $\square$

Let $G, K, \tau, \langle \cdot, \cdot \rangle$ be as at p. 4, and let M be the associated flag manifold. Then M is endowed with all the structures needed to define the Bourguignon-Li-Yau map, so for any probability measure γ on M there is a map $\Psi_\gamma : X := G/K \rightarrow \mathfrak{k}^*$ as in (66). Recall that $\mathcal{O} := \Phi(M) \subset \mathfrak{k}^*$ is a coadjoint orbit and that $\Phi : M \rightarrow \mathcal{O}$ is an equivariant symplectomorphism. Let $\widehat{\mathcal{O}}$ denote the convex hull of $\mathcal{O}$ and let Ω denote the interior of $\widehat{\mathcal{O}}$.

Lemma 69. (a) $0 \in \Omega$, (b) Ω is a nonempty open convex domain, (c) $\overline{\Omega} = \widehat{\mathcal{O}}$, (d) $\widehat{\mathcal{O}}$ is compact.

Proof. (a) For $v \in \mathfrak{k}$ set $h_v = \langle \Phi, v \rangle$. Since $\mathfrak{k} = [\mathfrak{k}, \mathfrak{k}]$, $v = \sum_{i=1}^k [u_i, w_i]$ for some $u_i, w_i \in \mathfrak{k}$, so by the equivariance of the moment map

$$h_v = \sum_{i=1}^k \langle \Phi, [u_i, w_i] \rangle = \sum_{i=1}^k \xi_{w_i} h_{u_i} = \sum_{i=1}^k \{h_{w_i}, h_{u_i}\}.$$

On the other hand, on a compact symplectic manifold Poisson brackets have zero mean:

$$\int_M \{f, g\} \omega^n = \int_M (X_f g) \omega^n = \int_M L_{X_f}(g \omega^n) = \int_M d i_{X_f}(g \omega^n) = 0.$$

Hence any component h_v of the moment map has zero mean. Therefore if $v \neq 0$ the function h_v necessarily changes sign. This shows that $\mathcal{O}$ is not contained in any halfspace with the origin on its boundary, which proves (a). (b) and (c) immediately follow. To prove (d) it is enough to observe that $\mathcal{O}$ is compact. $\square$

Definition 70. We say that a probability measure γ on M is τ -admissible if it does not charge the hyperplane sections of $M \subset \mathbb{P}(V)$, i.e. if $\gamma(H \cap M) = 0$ for any hyperplane $H \subset \mathbb{P}(V)$.

This condition is of course satisfied by any measure that is absolutely continuous with respect to the Riemannian measure. More generally, if $Z \subset M$ is a complex submanifold such that $Z \subset \mathbb{P}(V)$ is full (i.e. Z is not contained in any hyperplane) and γ is a smooth measure on Z , then γ , seen as a measure on M , is τ -admissible.

Lemma 71. Assume that $p_n \rightarrow p$ in $\overline{X}_\tau^S$ and let $\check{p}_n, \check{p}$ be as in (47). If γ is τ -admissible, then $\check{p}_n \rightarrow \check{p}$ γ -a.e.

Proof. By Lemma 14 (c) we can find unique $A_n, A \in \mathcal{H}(V)$ such that $\text{tr } A_n = \text{tr } A = 1$, $p_n = [A_n]$, $p = [A]$ and $A_n \rightarrow A$ in $\text{End } V$. Assume $p_n \in i_{W_n}(X_{W_n})$ and $p \in i_W(X_W)$. The set $N = \bigcup_n \mathbb{P}(W_n^\perp) \cup \mathbb{P}(W^\perp)$ has γ -measure zero since γ is τ -admissible. By Lemma 44 $\sqrt{A_n} \rightarrow \sqrt{A}$. If $x = [v] \in M - N$, then $\check{p}_n(x) = [\sqrt{A_n}v] \rightarrow [\sqrt{A}v] = \check{p}(x)$. This concludes the proof. $\square$

Theorem 72. For any τ -admissible probability measure γ on M the map Ψ_γ admits a continuous extension to $\overline{X}_\tau^S$ that we still denote by Ψ_γ . The extension is unique and satisfies $\Psi_\gamma(\overline{X}_\tau^S) \subset \widehat{\mathcal{O}}$. If $p \in X_W$, then

$$(73) \quad \Psi_\gamma(p) = \int_M \Phi(\check{p}(x)) d\gamma(x).$$

Proof. Let $p \in i_W(X_W)$. By Lemma 48 (a) the rational map $\check{p}$ is defined outside $\mathbb{P}(W^\perp)$. Since γ is τ -admissible $M - \mathbb{P}(W^\perp)$ has full measure. By Lemma 48 (e) $\check{p}(M - \mathbb{P}(W)) \subset M_W \subset M$, so the function $x \mapsto \Phi(\check{p}(x))$ is defined γ -a.e. on M and is bounded, hence integrable w.r.t. γ . This shows that formula (73) yields a well-defined map from $\overline{X}_\tau^S$ to $\mathfrak{k}^*$. If $W = V$ and $x = [v] \in M$, then $\check{p}([v]) = [\sqrt{\tau(g)\tau(g)^*}v] = \rho(g) \cdot x$. Therefore (73) agrees with (66) and yields an extension to $\overline{X}_\tau^S$ of the map previously defined on X . Let $\{p_n\} \subset \overline{X}_\tau^S$ be a sequence converging to some point p . By Lemma 71 $\Phi \circ \check{p}_n \rightarrow \Phi \circ \check{p}$ γ -a.e. Since Φ is bounded on M the integrals converge by the dominated convergence theorem. This shows that the map defined by (73) is continuous on $\overline{X}_\tau^S$. The uniqueness of a continuous extension is obvious since X is dense in $\overline{X}_\tau^S$. By Lemma 68 $\Psi_\gamma(X) \subset \widehat{\mathcal{O}}$, hence by continuity $\Psi_\gamma(\overline{X}_\tau^S) \subset \widehat{\mathcal{O}}$. $\square$

Now fix a τ -connected subspace $W \subsetneq V$. If ν is a probability measure on M_W denote by

$$\Psi_\nu^W : X_W \rightarrow \mathfrak{k}_W^*$$

the Bourguignon-Li-Yau map of M_W endowed with the structures induced from M .

Lemma 74. *For any $q \in X_W$,*

$$(75) \quad \Psi_\gamma(i_W(q)) = \Theta_W(\Psi_{\tilde{\gamma}}^W(q))$$

where $\tilde{\gamma} := (\tilde{\pi}_W)_\# \gamma$ is the push-forward probability measure on M_W and Θ_W is defined in (63). If γ is the K -invariant probability measure on M , then $\tilde{\gamma}$ is the K_W -invariant probability measure on M_W .

Proof. Assume $q = gK_W$ for $g \in S_W$. Then $p := i_W(q) = [\tau_W(g)\tau_W(g)^* \oplus 0] \in \overline{X}_\tau^S$. Recall from Lemma 48 (f) that for any $x \in M - \mathbb{P}(W)$ we have $\tilde{p}(x) = \rho(g) \cdot \tilde{\pi}_W(x)$. Therefore

$$(76) \quad \Psi_\gamma(p) = \int_M \Phi(\tilde{p}(x)) d\gamma(x) = \int_M \Phi(\rho(g) \cdot \tilde{\pi}_W(x)) d\gamma(x) = \int_{M_W} \Phi(\rho(g) \cdot y) d\tilde{\gamma}(y)$$

Using Lemma 64 we get

$$\begin{aligned} \Psi_\gamma(p) &= \int_{M_W} \Phi(\rho(g) \cdot y) d\tilde{\gamma}(y) = \int_{M_W} \Theta_W(\Phi^W(\rho(g) \cdot y)) d\tilde{\gamma}(y) = \\ &= \Theta_W\left(\int_{M_W} \Phi^W(\rho(g) \cdot y) d\tilde{\gamma}(y)\right) = \Theta_W(\Psi_{\tilde{\gamma}}^W(q)). \end{aligned}$$

This proves (75). To prove the last statement it is enough to take into account that the map $\tilde{\pi}_W$ is K_W -equivariant. $\square$

Remark 77. *If we use the scalar product $-B$ to identify $\mathfrak{k}^*$ with $\mathfrak{k}$, then formula (75) becomes simply $\Psi_\gamma(p) = \Psi_{\tilde{\gamma}}^W(p) + iZ_W$. This means that the restriction of Ψ_γ to any piece of the form $i_W(X_W)$ is just a translation (by iZ_W) of the Bourguignon-Li-Yau map on the corresponding M_W with respect to a suitable measure. In the case where γ is the K -invariant probability measure on M , this is particularly neat, since in that case $\tilde{\gamma}$ is just the K_W -invariant probability measure on M_W .*

Proposition 78. *Let W be a proper τ -connected subspace. Then $\Psi_\gamma(\overline{X}_\tau^S) \subset C_W$ and*

$$(79) \quad \Psi_\gamma^{-1}(E_W) = \bigsqcup_{W' \subset W} i_{W'}(X_{W'}) = \overline{i_W(X_W)}.$$

Proof. By Theorem 57 $\mathcal{O} \subset C_W$ so $\widehat{\mathcal{O}} \subset C_W$. By Theorem 72 $\Psi_\gamma(\overline{X}_\tau^S) \subset \widehat{\mathcal{O}}$ so $\Psi_\gamma(\overline{X}_\tau^S) \subset C_W$. This proves the first part of the theorem. Next we wish to show that $\Psi_\gamma(i_{W'}(X_{W'})) \subset E_W$ if $W' \subset W$. If $p \in i_{W'}(X_{W'})$, then $p = gK_{W'}$ for some $g \in S_{W'}$ and by (76)

$$\Psi_\gamma(p) = \int_{M_{W'}} \Phi(\rho(g) \cdot y) d\tilde{\gamma}(y).$$

Now, $\rho(g)M_{W'} \subset M_{W'} \subset M_W$, $\Phi(M_W) \subset E_W$ by Theorem 57 and E_W is obviously convex. Hence $\Psi_\gamma(p) \in E_W$. Conversely, fix $p \in i_{W'}(X_{W'})$ for some τ -connected subspace $W' \subset V$ and assume that $\Psi_\gamma(p) \in E_W$. We wish to prove that $W' \subset W$.

Fix a datum $(\mathfrak{h}, \Pi)$ and let x_0 be the line through the highest weight vector. Then by the definition of E_W we have

$$\begin{aligned} 0 &= \langle \Phi(x_0), v \rangle - \langle \Psi_\gamma(p), v \rangle = \\ &= \int_M \langle \Phi(x_0) - \Phi(\tilde{p}(x)), v \rangle d\gamma(x) = \int_M (h_v(x_0) - h_v(\tilde{p}(x))) d\gamma(x). \end{aligned}$$

Let $v_1, \dots, v_r$ be a basis of $\mathfrak{t}_I$ such that $v_j \in \mathfrak{c}_{\bar{W}, \mathfrak{h}, \Pi}^-$ for any j . By Theorem 57 $h_{v_j} \leq h_{v_j}(x_0)$, so $h_{v_j}(\tilde{p}(x)) = h_{v_j}(x_0)$ γ -a.e. for any j . For any j there is a subset $E_j \subset M$ such that $h_{v_j} \circ \tilde{p} = h_{v_j}(x_0)$ on $M - E_j$ and $\gamma(E_j) = 0$. Set

$$E := \cup_{j=1}^r E_j \cup \mathbb{P}(W'^\perp).$$

Then $\gamma(E) = 0$. By linearity $h_v \circ \tilde{p} = h_v(x_0)$ on $M - E$ for any $v \in \mathfrak{t}_I$. By Theorem 57 this means that $\tilde{p}(M - E) \subset M_W \subset \mathbb{P}(W)$. If $p = i_{W'}(gK_W)$, then $\tilde{p} = \rho(g) \circ \tilde{\pi}_{W'} = \tilde{\pi}_{W'} \circ \rho(g)$. Set $F = \rho(g)(E)$. Then $\tilde{\pi}_{W'}(M - F) = \tilde{p}(M - E) \subset \mathbb{P}(W)$. We claim that the projective subspace generated by $\tilde{\pi}_{W'}(M - F)$ coincides with $\mathbb{P}(W')$. Of course $\tilde{\pi}_{W'}(M - F) \subset \mathbb{P}(W')$, so the subspace generated is contained in $\mathbb{P}(W')$. On the other hand if $U \subset W'$ is a subspace such that $\tilde{\pi}_{W'}(M - F) \subset \mathbb{P}(U)$, then

$$\begin{aligned} M - F &\subset \mathbb{P}(U + (W')^\perp) \quad \rho(g)^{-1}F = E \\ M - E &= \rho(g)^{-1}(M - F) \subset \mathbb{P}((\rho(g))^{-1}(U + (W')^\perp)). \end{aligned}$$

Since $\gamma(M - E) = 1$ we get

$$\rho(g)^{-1}(U + (W')^\perp) = V \quad U + (W')^\perp = V.$$

Since $U \subset W'$ this yields $U = W'$. The claim is proved. Now from $\tilde{\pi}_{W'}(M - F) \subset \mathbb{P}(W)$ we get $\mathbb{P}(W') \subset \mathbb{P}(W)$ so $W' \subset W$. This proves the first equality in (79). The second one follows from Satake analysis of the boundary components. $\square$

It is useful to recall a few definitions and results regarding convex bodies (see e.g. [32]). If V is a real vector space and $E \subset V$ the *relative interior* of E , denoted $\text{relint } E$ is the interior of E in its affine hull. If E is a convex set in V a *face* of E is a convex subset $F \subset E$ with the following property: if $x, y \in E$ and $\text{relint}[x, y] \cap F \neq \emptyset$ then $[x, y] \subset F$. The *extreme points* of E are the points $x \in E$ such that $\{x\}$ is a face. If E is compact the faces are closed [32, p. 62]. If F is a face of E we say that $\text{relint } F$ is an *open face* of E .

Lemma 80. *Let V be a finite-dimensional real vector space and let $E \subset V$ be a compact convex set. Let $\{\lambda_\alpha\}_{\alpha \in A}$ be a family of linear functionals on V and let $a_\alpha \in \mathbb{R}$ be such that $E \subset \{v \in V : \lambda_\alpha(v) \leq a_\alpha \text{ for any } \alpha \in A\}$. Then*

$$F := \{v \in E : \lambda_\alpha(v) = a_\alpha \text{ for any } \alpha \in A\}$$

is empty or a face of E .

Proof. Fix $\alpha \in A$ and $x, y \in E$. If $\text{relint}[x, y] \cap F \neq \emptyset$, the affine function λ_α takes its maximum at an interior point of $[x, y]$. Therefore it must be constant on $[x, y]$. $\square$

Theorem 81 ([32, p. 62]). *If E is a compact convex set and F_1, F_2 are distinct faces of E then $\text{relint } F_1 \cap \text{relint } F_2 = \emptyset$. If G is a nonempty convex subset of E which is open in its affine hull, then $G \subset \text{relint } F$ for some face F of E . Therefore E is the disjoint union of its open faces.*

Lemma 82. *Let V be a finite-dimensional real vector space, $E \subset V$ a compact convex set and $A \subset E$ a convex set. Assume that A is dense in E and is open in its affine hull. Then $A = \text{relint } E$.*

Lemma 83. (a) *For any τ -connected subspace W the set $F_W := \widehat{\mathcal{O}} \cap E_W$ is a face of $\widehat{\mathcal{O}}$.*

- (b) *If W and W' are τ -connected subspaces of V and $F_W \subset F_{W'}$, then $W \subset W'$.*
(c) *$\text{relint } F_W = \text{relint } F_{W'}$ if and only if $W = W'$.*

Proof. (a) It follows from Theorem 57 that $\widehat{\mathcal{O}} \subset C_W$. By Lemma 55 E_W is the affine subspace where the inequalities defining C_W become equalities. Hence Lemma 80 ensures that F_W is a face of $\widehat{\mathcal{O}}$. (b) If $F_W \subset F_{W'}$ then $E_W \cap \mathcal{O} \subset E_{W'} \cap \mathcal{O}$. Using again Theorem 57 we get that

$$M_W = \Phi^{-1}(E_W) \subset \Phi^{-1}(E_{W'}) = M_{W'}.$$

By Corollary 32 this implies that $W \subset W'$. (c) If $\text{relint } F_W = \text{relint } F_{W'}$ also $F_W = F_{W'}$ by taking the closures. Hence $W = W'$ by (b). $\square$

Theorem 84. *Let γ be a τ -admissible measure. Then $\Psi_\gamma(\partial \overline{X}_\tau^S) \subset \partial \widehat{\mathcal{O}}$.*

Proof. By Theorems 24 and 36 the boundary of $\overline{X}_\tau^S$ is the disjoint union of $i_W(X_W)$ as W varies over all proper τ -connected subspaces. By Proposition 78 for any such W , $\Psi_\gamma(i_W(X_W)) \subset E_W \cap \widehat{\mathcal{O}} = F_W$ and by Lemma 83 F_W is a proper face of $\widehat{\mathcal{O}}$, so $F_W \subset \partial \widehat{\mathcal{O}}$. $\square$

3.2. The case of the K -invariant measure. Let $G, K, \tau, \langle \cdot, \cdot \rangle$ be as at p. 4, and let M be the associated flag manifold. Let μ denote the unique K -invariant probability measure on M , i.e. the Riemannian volume normalized to be of unit mass. Set

$$\Psi := \Psi_\mu.$$

Theorem 85. $\Psi|_X$ is a diffeomorphism of X onto the interior of $\widehat{\mathcal{O}}$.

Proof. Consider the map

$$F : G \rightarrow \mathfrak{k}^* \quad F(g) := \int_M \Phi(g \cdot x) d\mu(x).$$

Fix $g \in G$, $v \in \mathfrak{k}$ and set $\tilde{\mu} = g_\# \mu$. Denote by $R_g : G \rightarrow G$ the right translation and set $w := (dR_g)(e)(iv) \in T_g G$.

$$\begin{aligned} dF(g)(w) &= \left. \frac{d}{dt} \right|_{t=0} F(\exp(itv)g) = \left. \frac{d}{dt} \right|_{t=0} \int_M \Phi(\exp(itv) \cdot y) d\tilde{\mu}(y) \\ \langle dF(g)(w), v \rangle &= \int_M \left. \frac{d}{dt} \right|_{t=0} \langle \Phi(\exp(itv) \cdot y), v \rangle d\tilde{\mu}(y) \\ \left. \frac{d}{dt} \right|_{t=0} \langle \Phi(\exp(itv) \cdot y), v \rangle &= d\langle \Phi, v \rangle(y)(\xi_{iv}(y)) = -i_{\xi_v} \omega(J\xi_v(y)) = -|\xi_v(y)|^2 \\ \langle dF(g)(w), v \rangle &= - \int_M |\xi_v|^2 d\tilde{\mu}. \end{aligned}$$

Since $\tilde{\mu}$ is a smooth measure with strictly positive density, if $dF(g)(w) = 0$, then $\xi_v \equiv 0$ and $v = 0$. This shows that the restriction of $dF(g)$ to $(dR_g)(e)(i\mathfrak{k}) \subset T_g G$ is

an isomorphism. In particular $dF(g)$ is onto and F is a submersion. Let $g = \rho(g)a$ be the polar decomposition, i.e. $a \in K$. Since $a_{\#}\mu = \mu$

$$\begin{aligned} F(g) &:= \int_M \Phi(g \cdot x) d\mu(x) = \int_M \Phi(\rho(g) \cdot x) d(a_{\#}\mu)(x) = \int_M \Phi(\rho(g) \cdot x) d\mu(x) = \\ &= \Psi(gK). \end{aligned}$$

So if $\pi : G \rightarrow X$ is the canonical map $\pi(g) = gK$, then $\Psi|_X \circ \pi = F$. Since F is a submersion, $\Psi|_X$ is a local diffeomorphism and in particular an open map. So $\Psi(X)$ is open in $\mathfrak{k}^*$. Since $\Psi(X) \subset \widehat{\mathcal{O}}$ by Lemma 68, it follows that $\Psi(X) \subset \text{int } \widehat{\mathcal{O}} = \Omega$. On the other hand, by Theorem 84, $\Psi(\partial \overline{X}_\tau^S) \subset \partial \widehat{\mathcal{O}}$. Therefore $\Psi|_X : X \rightarrow \Omega$ is also a proper map. It follows that it is a covering of Ω and therefore a diffeomorphism. $\square$

Theorem 86. (a) *The Bourguignon-Li-Yau map Ψ is a homeomorphism of $\overline{X}_\tau^S$ onto $\widehat{\mathcal{O}}$.*
 (b) *For any τ -connected subspace W , $\Psi(i_W(X_W)) = \text{relint } F_W$ and $\Psi \circ i_W$ is a diffeomorphism of X_W onto $\text{relint } F_W$.*

Proof. By Theorem 72 the map $\Psi : \overline{X}_\tau^S \rightarrow \mathfrak{k}^*$ is continuous and its image is contained in $\widehat{\mathcal{O}}$. By Theorem 84 $\Psi(\partial \overline{X}_\tau^S) \subset \partial \widehat{\mathcal{O}}$. Since $\overline{X}_\tau^S$ is compact and $\Psi(\overline{X}_\tau^S) \supset \Psi(X) = \Omega$, $\Psi(\overline{X}_\tau^S) = \widehat{\mathcal{O}}$, so Ψ is surjective. Next consider a proper τ -connected subspace W . By Proposition 78 $\Psi(i_W(X_W)) \subset \widehat{\mathcal{O}} \cap E_W$. By Lemma 74 the diagram

$$(87) \quad \begin{array}{ccc} X_W & \xrightarrow{\Psi^W} & \mathfrak{k}_W^* \\ i_W \downarrow & & \downarrow \Theta_W \\ i_W(X_W) & \xrightarrow{\Psi} & \mathfrak{k}^* \end{array}$$

commutes. (Here Ψ^W is the Bourguignon-Li-Yau map of M_W with the invariant probability measure.) Set $\mathcal{O}_W := \Phi^W(M_W) \subset \mathfrak{k}_W^*$, $\widehat{\mathcal{O}}_W := \text{convex hull of } \mathcal{O}_W$ and $\Omega_W := \text{int } \widehat{\mathcal{O}}_W$. By what we have just proved applied to the flag manifold M_W , $\Psi^W(\overline{X}_W) = \widehat{\mathcal{O}}_W$ and $\Psi^W|_{X_W}$ is a diffeomorphism onto Ω_W . Since Θ_W is an affine embedding, it follows that $\Psi \circ i_W$ is a diffeomorphism of X_W onto $\Theta_W(\Omega_W)$. We wish to show that

$$(88) \quad \Theta_W(\Omega_W) = \text{relint} \left(E_W \cap \widehat{\mathcal{O}} \right).$$

Let $\overline{X}_W$ denote the Satake compactification of X_W associated to τ_W and let $\overline{i_W(X_W)}$ denote the closure of $i_W(X_W)$ in $\overline{X}_\tau^S$. If in (87) we replace X_W by $\overline{X}_W$ and $i_W(X_W)$ by $\overline{i_W(X_W)}$, then by continuity the diagram still commutes. Since Ψ is surjective, Proposition 78 implies that $\Psi(\overline{i_W(X_W)}) = E_W \cap \widehat{\mathcal{O}}$, hence

$$\Theta_W \circ \Psi^W(\overline{X}_W) = \Psi(\overline{i_W(X_W)}) = E_W \cap \widehat{\mathcal{O}}.$$

So $\Theta_W(\Omega_W) = \Theta_W \circ \Psi^W(X_W)$ is dense in $E_W \cap \widehat{\mathcal{O}}$. As Ω_W is open in $\mathfrak{k}_W^*$, $\Theta_W(\Omega_W)$, is an open convex domain in $\Theta_W(\mathfrak{k}_W^*)$ which is its affine hull. Lemma 82 implies that (88) holds true and concludes the proof of (b). Now we wish to prove that Ψ is injective on the boundary. By the above it is injective on X and also on each boundary component $i_W(X_W)$. Moreover each boundary component $i_W(X_W)$ is mapped to $\text{relint } F_W$. If $i_W(X_W) \neq i_{W'}(X_{W'})$, then $W \neq W'$ hence, by Lemma

83 (c), $\text{relint } F_W \neq \text{relint } F_{W'}$. But F_W and $F_{W'}$ are faces of $\widehat{\mathcal{O}}$, by Lemma 83 (a), hence $\text{relint } F_W \cap \text{relint } F_{W'} = \emptyset$. It follows that Ψ is injective. Since $\overline{X}_\tau^S$ is compact and $\widehat{\mathcal{O}}$ is Hausdorff Ψ is a homeomorphism. This finally proves (a). $\square$

Theorem 89. *Let γ be a τ -admissible measure. Then the Bourguignon-Li-Yau map is $\Psi_\gamma : \overline{X}_\tau^S \rightarrow \widehat{\mathcal{O}}$ is surjective.*

Proof. Set $\gamma_t := (1-t)\mu + t\gamma$ and

$$H : \overline{X}_\tau^S \times [0, 1] \rightarrow \widehat{\mathcal{O}} \quad H(p, t) := \Psi_{\gamma_t}(p).$$

γ_t is τ -admissible for every $t \in [0, 1]$, H is continuous and $H(\partial \overline{X}_\tau^S \times [0, 1]) \subset \partial \widehat{\mathcal{O}}$ by Theorem 84. Since $\Psi|_{\partial \overline{X}_\tau^S} = H(\cdot, 0)|_{\partial \overline{X}_\tau^S}$ is a homeomorphism, it has degree 1. Hence the same holds for $H(\cdot, 1)|_{\partial \overline{X}_\tau^S} = \Psi_\gamma|_{\partial \overline{X}_\tau^S}$. By a classical topological argument this yields the surjectivity of $H(\cdot, 1) = \Psi_\gamma$. $\square$

3.3. Furstenberg compactifications. Another way to compactify $X = G/K$ was found by Furstenberg [13] in his search for an analogue of the Poisson formula for the unit disc. We recall very briefly the definition in the case of type IV symmetric spaces (see [9, §I.6] for the general case). Let G be a connected complex semisimple Lie group and K a maximal compact subgroup. A homogeneous space $M = G/P$ is called a *Furstenberg boundary* of G if for every probability measure ν on M , there exists a sequence $g_j \in G$ and a point $x \in M$ such that $g_{j\#}\nu \rightarrow \delta_x$. Using Iwasawa structure theory, Moore [28, Thm. 1] proved that G/P is a boundary if and only if P is parabolic. In this case K acts transitively and M has a unique K -invariant probability measure μ . For any topological space Z , let $\mathcal{P}(Z)$ denote the set of Borel probability measures on Z provided with the weak topology. Since μ is K -invariant the map

$$G \rightarrow \mathcal{P}(M) \quad g \mapsto g_{\#}\mu$$

descends to a continuous map $i_M : X = G/K \rightarrow \mathcal{P}(M)$, which is injective iff P does not contain simple factors of G (see [28, Thm. 4] or [9, Prop. I.6.16]). In this case M is called a *faithful* Furstenberg boundary and the set

$$(90) \quad \overline{X}_M^F := \overline{i_M(X)}$$

is called the *Furstenberg compactification* of X associated to the faithful boundary M . Fix an irreducible complex representation $\tau : G \rightarrow \text{GL}(V)$ such that P is the stabilizer of some $x_0 \in \mathbb{P}(V)$ and $\ker \tau$ is finite. Such representations always exist. Then M can be identified with the orbit $G \cdot x_0$, which is the unique closed orbit in $\mathbb{P}(V)$. If a K -invariant Hermitian product is fixed on V , M becomes a flag manifold, so we are back to the previous setting.

Theorem 91. *The map*

$$\Gamma : \overline{X}_\tau^S \rightarrow \overline{X}_M^F \quad \Gamma(p) := \check{p}_{\#}\mu$$

is a G -equivariant homeomorphism of $\overline{X}_\tau^S$ onto $\overline{X}_M^F$ such that $i_M = \Gamma \circ i_\tau$ (compare (16)).

Proof. For any $p \in \overline{X}_\tau^S$ the rational map $\check{p} : M \dashrightarrow M$ is defined μ -a.e., since μ does not charge linear sections of $M \subset \mathbb{P}(V)$. Therefore $\Gamma(p) = \check{p}_{\#}\mu$ is well-defined

for any $p \in \overline{X}_\tau^S$. If $p_n \rightarrow p$, then $\check{p}_n \rightarrow \check{p}$ μ -a.e. by Lemma 71. It follows that $\Gamma(p_n) \rightarrow \Gamma(p)$. So Γ is continuous. Moreover $\Gamma \circ i_\tau = i_M$ (see (16) for the definition of i_τ). Indeed by Lemma 48 (f), if $p = i_\tau(gK)$, then $\check{p} = L_{\rho(g)} : M \rightarrow M$ so $\Gamma(p) = \rho(g)_\# \mu = g_\# \mu = i_M(gK)$. Therefore $\Gamma(M) = i_M(X)$ is dense in $\overline{X}_M^F$. Since $\overline{X}_\tau^S$ is compact it follows that Γ is surjective. Consider now the following maps

$$\begin{aligned} F : \mathcal{P}(M) &\rightarrow \mathcal{P}(\widehat{\mathcal{O}}) & F(\gamma) &= \Phi_\# \gamma \\ \mathbb{B} : \mathcal{P}(\widehat{\mathcal{O}}) &\rightarrow \widehat{\mathcal{O}} & \mathbb{B}(\nu) &= \int_{\widehat{\mathcal{O}}} z d\nu(z). \end{aligned}$$

$\mathbb{B}(\nu)$ is just the barycenter of ν . Now $\mathbb{B} \circ F \circ \Gamma = \Psi$. Since Ψ is a homeomorphism, Γ is injective. As $\mathcal{P}(M)$ is a Hausdorff space, Γ is a homeomorphism. Moreover $i_M = \Gamma \circ i_\tau$, so Γ is G -equivariant on X . By continuity it is equivariant also on the compactifications. $\square$

4. APPLICATION TO EIGENVALUE ESTIMATES

In this section we apply Theorem 89 to a problem in spectral geometry. Let M be a complex manifold and g a Kähler metric. Denote by $\Delta_g : C^\infty(M) \rightarrow C^\infty(M)$ the Laplace-Beltrami operator

$$\Delta_g f = -d^* df = 2 g^{i\bar{j}} \frac{\partial^2 f}{\partial z^i \partial \bar{z}^j}.$$

It is well-known that $-\Delta_g$ is a positive elliptic operator with eigenvalues $0 \leq \lambda_1(g) \leq \lambda_2(g) \leq \dots$. Assume that M is a Fano manifold and g_{KE} is a Kähler-Einstein metric with Kähler class $\omega_{KE} \in 2\pi c_1(M)$ (i.e. $\text{Ric}(g_{KE}) = g_{KE}$). Denote by $\mathfrak{k}$ the Lie algebra of Killing vector fields and by $\mathfrak{g}$ the algebra of holomorphic vector fields (considered as real fields). The map $v \mapsto Jv$ endows $\mathfrak{g}$ with the structure of complex Lie algebra. For $u \in C^\infty(M, \mathbb{R})$ let X_u denote the Hamiltonian vector field such that $du = -i_{X_u} \omega_{KE}$.

Theorem 92. *Let M be a Fano manifold and let g_{KE} be a Kähler-Einstein metric as above. Then (a) $\mathfrak{k}$ is a real form of $\mathfrak{g}$; (b) if $\mathfrak{g} \neq \{0\}$ then $\lambda_1(g_{KE}) = 2$. (c) Set $\Lambda_1 = \{u \in C^\infty(M, \mathbb{R}) : -\Delta_{KE} u = 2u\}$. If $u \in \Lambda_1$, then $X_u \in \mathfrak{k}$. (d) Define $L : \Lambda_1 \rightarrow \mathfrak{k}$ by $L(u) = X_u$. Then L is an isomorphism and the map $\Phi : M \rightarrow \mathfrak{k}^*$ defined by*

$$\langle \Phi(x), v \rangle := L^{-1}(v)(x)$$

is a moment map for the action of $\text{Isom}(M, g_{KE})$.

This statement combines results of Matsushima and Futaki, see [24, p. 95], [15, Thm. 2.4.3, p.41], [14, Lemma 4.2].

Assume now that M is a Hermitian symmetric space of the compact type. We are interested in upper bounds for $\lambda_1(g)$ that hold for any Kähler metric in $2\pi c_1(M)$. Using the notation (1) of the introduction, we wish to estimate $I(2\pi c_1(M))$. Let g_{KE} be the symmetric Kähler-Einstein metric with $\text{Ric}(g_{KE}) = g_{KE}$ and let ω_{KE} be its Kähler form. Let K denote the connected component of the identity of $\text{Isom}(M, g_{KE})$. K acts transitively on M . Set $n := \dim_{\mathbb{C}} M$.

Lemma 93. *Let $e_1, \dots, e_l$ be an orthonormal basis of $\mathfrak{k} := \text{Lie } K$ with respect to the scalar product $-B/2$ (B is the Killing form). Set $f_j = \langle \Phi, e_j \rangle$ and*

$$\alpha := \sum_{j=1}^l i \partial f_j \wedge \bar{\partial} f_j.$$

Then $\alpha = \omega_{KE}$ and $|\Phi|^2 = n$.

Proof. Fix a point $x_0 \in M$ and let K_0 be its stabilizer. If $\mathfrak{k} = \mathfrak{k}_0 \oplus \mathfrak{m}$, then $\mathfrak{m}$ is $\text{ad } K_0$ -invariant and we identify it with $T_{x_0}M$ via the mapping $v \mapsto \xi_v(x_0)$. By [25, p. 261] there is $H \in \mathfrak{z}(\mathfrak{k}_0)$ such that $J_{x_0} = \text{ad } H|_{\mathfrak{m}}$. The Kähler-Einstein metric g_{KE} is the one associated with the scalar product $h = -\frac{1}{2}B$ on $\mathfrak{m}$, see [25, p. 262]. Define a map $\Phi : M \rightarrow \mathfrak{k}^*$ by

$$(94) \quad \langle \Phi(a \cdot x_0), v \rangle = \frac{1}{2}B(H, \text{Ad}(a^{-1})v).$$

This map is well defined since $H \in \mathfrak{z}(K_0)$. Moreover Φ is K -equivariant. We claim that Φ is the moment map of (M, ω_{KE}) . Indeed for $w \in T_{x_0}M$

$$(95) \quad \begin{aligned} d\langle \Phi, v \rangle(x_0)(w) &= \left. \frac{d}{dt} \right|_{t=0} \langle \Phi(e^{tw} \cdot x_0), v \rangle = \left. \frac{1}{2} \frac{d}{dt} \right|_{t=0} B(H, \text{Ad}(e^{-tw})v) = \\ &= -\frac{1}{2}B(H, [w, v]) = -\frac{1}{2}B([v, H], w). \end{aligned}$$

Let $v = v_1 + v_2$ with $v_1 \in \mathfrak{k}_0$, $v_2 \in \mathfrak{m}$. Then $[v, H] = [v_2, H]$ and $\xi_v(x_0) = v_2$, so

$$\begin{aligned} d\langle \Phi, v \rangle(x_0)(w) &= \frac{1}{2}B([H, v_2], w) = -g_{KE}(J_{x_0}v_2, w) = -\omega_{KE}(\xi_{v_2}(x_0), w) = \\ &= -i_{\xi_v} \omega_{KE}(w). \end{aligned}$$

This shows that $d\langle \Phi, v \rangle = -i_{\xi_v} \omega_{KE}$ at x_0 . By K -equivariance this holds on all of M , so Φ is the moment map. A simple computation shows that for $f \in C^\infty(M, \mathbb{R})$ and tangent vectors X, Y

$$i\partial f \wedge \bar{\partial} f(X, Y) = \frac{1}{2}(Yf \cdot JXf - Xf \cdot JYf).$$

Apply this to $X = w_1, Y = w_2$ and $f = \langle \Phi, v \rangle$, where $w_i \in \mathfrak{m}$ and $v \in \mathfrak{k}$. Using (95) and $[H, [H, w_i]] = J^2 w_i = -w_i$ we get

$$\begin{aligned} i\partial f \wedge \bar{\partial} f(w_1, w_2) &= \\ &= \frac{1}{2} \left(\frac{1}{2}B(H, [v, w_2]) \frac{1}{2}B(H, [v, [H, w_1]]) - \frac{1}{2}B(H, [v, w_1]) \frac{1}{2}B(H, [v, [H, w_2]]) \right) = \\ &= \frac{1}{2} (h([H, w_2], v)h(v, [H, [H, w_1]]) - h([H, w_1], v)h(v, [H, [H, w_2]])) = \\ &= \frac{1}{2} (h([H, w_1], v)h(v, w_2) - h([H, w_2], v)h(v, w_1)). \end{aligned}$$

Since e_j is an orthonormal basis of $(\mathfrak{k}, h)$

$$\begin{aligned} \alpha(w_1, w_2) &= \frac{1}{2} \sum_{j=1}^l (h([H, w_1], e_j)h(e_j, w_2) - h([H, w_2], e_j)h(e_j, w_1)) = \\ &= \frac{1}{2} (g_{KE}(Jw_1, w_2) - g_{KE}(Jw_2, w_1)) = \omega_{KE}(w_1, w_2). \end{aligned}$$

This proves that $\alpha = \omega_{KE}$. The norm $|\Phi|$ is constant since K is transitive. Using (94) we get

$$\begin{aligned} |\Phi|^2(x_0) &= \sum_{j=1}^l \langle \Phi, e_j \rangle^2 = \sum_j \left[\frac{1}{2} B(H, e_j) \right]^2 = \sum_j [h(H, e_j)]^2 = \\ &= h(H, H) = -\frac{1}{2} B(H, H). \end{aligned}$$

But $(\text{ad}H)^2$ is 0 on $\mathfrak{k}_0$ and is -1 on $\mathfrak{m}$. Since $\dim_{\mathbb{R}} \mathfrak{m} = 2n$, $B(H, H) = -2n$ and $|\Phi|^2 = n$. $\square$

Theorem 96. *Let M be a Hermitian symmetric space of the compact type. Let N be a Kähler manifold and let $a \in H^2(N)$ be a Kähler class. Let $F : N \rightarrow M$ be a holomorphic map and assume that no nontrivial section of $-K_M$ vanishes on $F(N)$. Let $n = \dim_{\mathbb{C}} M$ and $d = \dim_{\mathbb{C}} N$. Then for any Kähler metric g with Kähler form $\omega \in a$*

$$(97) \quad \lambda_1(g) \leq \frac{4\pi d}{n} \cdot \frac{\int_N F^*(c_1(M)) \cup a^{d-1}}{\int_N a^d}.$$

Proof. Let g_{KE} , ω_{KE} , K , $\mathfrak{k}$ and $\mathfrak{g}$ have the same meaning as above. Let G denote the connected component of the identity of the automorphism group of M . By Theorem 92 K is a maximal compact subgroup of G . Set $X := G/K$. Since K acts transitively the norm $|\Phi|$ is constant. Therefore, using the notation of Lemma 93 and setting $l := \dim K$,

$$\sum_{j=1}^l \int_M f_j^2 \frac{\omega_{KE}^n}{n!} = |\Phi|^2 \text{vol}(M, g_{KE}).$$

The anticanonical bundle $-K_M$ is very ample. Set $V := H^0(M, -K_M)^*$. Since $-K_M$ is G -equivariant there is an obvious representation $\tau : G \rightarrow \text{GL}(V)$. The metric g_{KE} induces an L^2 -scalar product $\langle \cdot, \cdot \rangle$ on V which is K -invariant. By the Borel-Weil theorem the representation τ is irreducible and the map $\varphi : M \rightarrow \mathbb{P}(V)$ induced by the linear system $| -K_M |$ is a biholomorphism onto the unique closed orbit in $\mathbb{P}(V)$. Let g_{FS} be the Fubini-Study metric on $\mathbb{P}(V)$ induced by $\langle \cdot, \cdot \rangle$, and let ω_{FS} be its Kähler form. Since M is a compact symmetric space any cohomology class contains a unique invariant form [36, Thm. 8.5.8 p. 250]. Hence $\varphi^* \omega_{FS} = \omega_{KE}$. So the data $G, K, \tau, \langle \cdot, \cdot \rangle$ satisfy the same assumption set forth at p. 4 and (M, ω_{KE}) is the associated flag manifold as defined in §2.1. Let $\Phi : M \rightarrow \mathfrak{k}^*$ be the moment map of M provided with these structures, set $\mathcal{O} = \Phi(M)$ and let Ω be the interior of $\widehat{\mathcal{O}}$. Set

$$\gamma := \frac{1}{\text{vol}(N, g)} F_{\#} \text{vol}_g.$$

Any hyperplane $H \subset \mathbb{P}(V)$ is defined by the vanishing of a section $s \in V^*$, so by hypothesis $F^{-1}(H)$ is a proper complex subvariety of N . Therefore γ is a τ -admissible measure on M . By Theorem 89 $\Psi_{\gamma}(X) = \Omega$ and by Lemma 69 $0 \in \Omega$. This means that there some $b \in G$ such that

$$(98) \quad \Psi_{\gamma}(bK) = \int_M \Phi(\rho(b) \cdot x) d\gamma(x) = 0.$$

If we set $a = \rho(b)$ and $h_j = f_j \circ a \circ F$, then the above formula yields

$$\int_N h_j(y) \operatorname{vol}_g(y) = 0$$

for $j = 1, \dots, l$. So the functions h_j have zero mean. By the Rayleigh theorem we get

$$(99) \quad \lambda_1(N, g) \leq \frac{\sum_{j=1}^l \int_N |\nabla h_j|_g^2 \frac{\omega^d}{d!}}{\sum_{j=1}^l \int_N h_j^2 \frac{\omega^d}{d!}}.$$

We have to estimate the right hand side. Since K is transitive, the norm $|\Phi|$ is constant. By Lemma 93

$$(100) \quad \sum_{j=1}^l \int_N h_j^2 \frac{\omega^d}{d!} = |\Phi|^2 \operatorname{vol}(N, g) = \frac{n}{d!} \int_N a^d.$$

Recall that for any $u \in C^\infty(N, \mathbb{R})$

$$i\partial u \wedge \bar{\partial} u \wedge \frac{\omega^{d-1}}{(d-1)!} = \frac{1}{2} |\nabla u|^2 \frac{\omega^d}{d!}.$$

Hence

$$\begin{aligned} \sum_{j=1}^l \int_N |\nabla h_j|_g^2 \frac{\omega^d}{d!} &= \frac{2}{(d-1)!} \sum_{j=1}^l \int_N i\partial h_j \wedge \bar{\partial} h_j \wedge \omega^{d-1} = \\ &= \frac{2}{(d-1)!} \int_N F^*(a^* \alpha) \wedge \omega^{d-1}. \end{aligned}$$

Since α is closed and G is connected the form $a^* \alpha$ is cohomologous to α . Therefore

$$\sum_{j=1}^l \int_N |\nabla h_j|_g^2 \frac{\omega^d}{d!} = \frac{4\pi}{(d-1)!} \int_N F^*(c_1(M)) \cup a^{d-1}.$$

Using (100) we get the result. $\square$

Remark 101. We wish to compare this result with those contained in [2] which deal with a similar situation in the case of the Grassmannian. Let V be a vector space of dimension k . Denote by M the Grassmannian of r -planes in V , by $U \rightarrow M$ the tautological bundle ($U_x = x$) and by $\mathcal{O}_M(1)$ the hyperplane bundle of the Plücker embedding. Then $V^* = H^0(M, U^*)$, $\Lambda^r U^* = \mathcal{O}_M(1)$, $-K_M = \mathcal{O}_M(k)$ and $H^0(M, \mathcal{O}_M(1)) = \Lambda^r V^*$.

Assume that $E \rightarrow N$ is a globally generated holomorphic vector bundle of rank r with $H^0(N, E) = V^*$. Consider the map $F : N \rightarrow M$, $F(x) = \operatorname{Ann}(\{s \in V^* : s(x) = 0\})$. By construction $E = F^*(U^*)$. We claim that if the map F satisfies the condition of Theorem 96, i.e. no section of $-K_M$ vanishes on $F(N)$, then the Gieseker point T_E is stable. Indeed, since $-K_M = \mathcal{O}_M(k)$, no section of $\mathcal{O}_M(1)$ vanishes on $F(N)$, hence the map $F^* : H^0(M, \mathcal{O}_M(1)) = H^0(M, \Lambda^r U^*) \rightarrow H^0(N, \Lambda^r E)$ is injective and the Gieseker points of E and U^* are related by $T_E = F^* \circ T_{U^*}$. It follows that T_E is stable. So in this particular case, where F is the map corresponding to the full linear system of sections of E , our Theorem 96 is a consequence of Theorem 1.1 in [2].

More generally, assume that N is a manifold, M is the Grassmannian of r -planes in some V and $F : N \rightarrow M$ is some map satisfying the assumption of Theorem 96.

Since $F^* : H^0(M, \mathcal{O}_M(1)) = \Lambda^r V \rightarrow H^0(N, E)$ is injective and Λ^r commutes with pull-back, also $F^* : H^0(M, U^*) = V \rightarrow H^0(N, E)$ is injective. The constructions in the proof of Theorem 96 relate to $V = H^0(M, U)$ or, equivalently, to the subspace $\text{Im } F^* \subset H^0(N, E)$. Indeed coupled with a theorem of Xiaowei Wang (see [35, Thm. 3.1], [7, Thm. 2.5]) formula (98) says roughly that $\text{Im } F^*$ is " ω -balanced" (see [2, p. 380]). This does not imply that the whole space $H^0(N, E)$ be ω -balanced (which is equivalent to T_E being stable). So for general F the hypothesis of Theorem 96 is weaker than the assumption of Theorem 1.1 in [2].

Corollary 102. *Let M be a Hermitian symmetric space of compact type. If g is a Kähler metric with Kähler form $\omega \in 2\pi c_1(M)$, then*

$$\lambda_1(M, g) \leq 2.$$

The bound is attained by the symmetric metric.

Proof. To get the estimate it is enough to apply the theorem with $N = M$, $F = \text{Id}$ and $a = 2\pi c_1(M)$. The bound is attained by g_{KE} by Theorem 92. $\square$

REFERENCES

- [1] D. N. Akhiezer. *Lie group actions in complex analysis*. Aspects of Mathematics, E27. Friedr. Vieweg & Sohn, Braunschweig, 1995.
- [2] C. Arezzo, A. Ghigi, and A. Loi. Stable bundles and the first eigenvalue of the Laplacian. *J. Geom. Anal.*, 17(3):375–386, 2007.
- [3] M. F. Atiyah. Convexity and commuting Hamiltonians. *Bull. London Math. Soc.*, 14(1):1–15, 1982.
- [4] M. Audin. *Torus actions on symplectic manifolds*, volume 93 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, revised edition, 2004.
- [5] R. J. Baston and M. G. Eastwood. *The Penrose transform*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, 1989.
- [6] M. Berger. Sur les premières valeurs propres des variétés riemanniennes. *Compositio Math.*, 26:129–149, 1973.
- [7] L. Biliotti and A. Ghigi. Homogeneous bundles and the first eigenvalue of symmetric spaces. *Ann. Inst. Fourier (Grenoble)*, 58(7):2315–2331, 2008.
- [8] D. D. Bleecker. The spectrum of a Riemannian manifold with a unit Killing vector field. *Trans. Amer. Math. Soc.*, 275(1):409–416, 1983.
- [9] A. Borel and L. Ji. *Compactifications of symmetric and locally symmetric spaces*. Mathematics: Theory & Applications. Birkhäuser Boston Inc., Boston, MA, 2006.
- [10] J.-P. Bourguignon, P. Li, and S.-T. Yau. Upper bound for the first eigenvalue of algebraic submanifolds. *Comment. Math. Helv.*, 69(2):199–207, 1994.
- [11] B. Colbois and J. Dodziuk. Riemannian metrics with large λ_1 . *Proc. Amer. Math. Soc.*, 122(3):905–906, 1994.
- [12] Y. Colin de Verdière. Construction de laplaciens dont une partie finie du spectre est donnée. *Ann. Sci. École Norm. Sup. (4)*, 20(4):599–615, 1987.
- [13] H. Furstenberg. A Poisson formula for semi-simple Lie groups. *Ann. of Math. (2)*, 77:335–386, 1963.
- [14] A. Futaki. The Ricci curvature of symplectic quotients of Fano manifolds. *Tohoku Math. J. (2)*, 39(3):329–339, 1987.
- [15] A. Futaki. *Kähler-Einstein metrics and integral invariants*. Springer-Verlag, Berlin, 1988.
- [16] R. Goodman and N. R. Wallach. *Symmetry, representations, and invariants*, volume 255 of *Graduate Texts in Mathematics*. Springer, Dordrecht, 2009.
- [17] V. Guillemin and S. Sternberg. *Symplectic techniques in physics*. Cambridge University Press, Cambridge, second edition, 1990.
- [18] Y. Guivarc’h, L. Ji, and J. C. Taylor. *Compactifications of symmetric spaces*, volume 156 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1998.
- [19] P. R. Halmos. *Measure Theory*. D. Van Nostrand Company, Inc., New York, N. Y., 1950.

- [20] J. Hersch. Quatre propriétés isopérimétriques de membranes sphériques homogènes. *C. R. Acad. Sci. Paris Sér. A-B*, 270:A1645–A1648, 1970.
- [21] A. Huckleberry. Introduction to group actions in symplectic and complex geometry. In *Infinite dimensional Kähler manifolds (Oberwolfach, 1995)*, volume 31 of *DMV Sem.*, pages 1–129. Birkhäuser, Basel, 2001.
- [22] A. W. Knap. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 2002.
- [23] A. W. Knap. *Advanced real analysis*. Cornerstones. Birkhäuser Boston Inc., Boston, MA, 2005.
- [24] S. Kobayashi. *Transformation groups in differential geometry*. Springer-Verlag, New York, 1972. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 70.
- [25] S. Kobayashi and K. Nomizu. *Foundations of differential geometry. Vol I*. Interscience Publishers, a division of John Wiley & Sons, New York-London, 1963.
- [26] A. Korányi. Remarks on the Satake compactifications. *Pure Appl. Math. Q.*, 1(4, part 3):851–866, 2005.
- [27] D. McDuff and D. Salamon. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1998.
- [28] C. C. Moore. Compactifications of symmetric spaces. *Amer. J. Math.*, 86:201–218, 1964.
- [29] G. Ottaviani. Rational homogeneous varieties. Notes from a course held in Cortona, Italy, 1995. <http://www.math.unifi.it/~ottaviani/public.html>.
- [30] R. Sanyal, F. Sottile, and B. Sturmfels. Orbitopes. *arXiv:math.AG/0911.5436*, 2009. Preprint.
- [31] I. Satake. On representations and compactifications of symmetric Riemannian spaces. *Ann. of Math. (2)*, 71:77–110, 1960.
- [32] R. Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1993.
- [33] J.-P. Serre. Représentations linéaires et espaces homogènes kähleriens des groupes de Lie compacts (d’après Armand Borel et André Weil). In *Séminaire Bourbaki, Vol. 2*, pages Exp. No. 100, 447–454. Soc. Math. France, Paris, 1995.
- [34] È. B. Vinberg, editor. *Lie groups and Lie algebras, III*. Springer-Verlag, Berlin, 1994. Structure of Lie groups and Lie algebras, A translation of *Current problems in mathematics. Fundamental directions. Vol. 41* (Russian), Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1990 [MR 91b:22001], Translation by V. Minachin [V. V. Minakhin], Translation edited by A. L. Onishchik and È. B. Vinberg.
- [35] X. Wang. Balance point and stability of vector bundles over a projective manifold. *Math. Res. Lett.*, 9(2-3):393–411, 2002.
- [36] J. A. Wolf. *Spaces of constant curvature*. McGraw-Hill Book Co., New York, 1967.
- [37] P. C. Yang and S. T. Yau. Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 7(1):55–63, 1980.
- [38] F. Ziegler. On the Kostant convexity theorem. *Proc. Amer. Math. Soc.*, 115(4):1111–1113, 1992.

UNIVERSITÀ DI PARMA

E-mail address: leonardo.biliotti@unipr.it

UNIVERSITÀ DI MILANO BICOCCA

E-mail address: alessandro.ghigi@unimib.it